

## Exploration

# From Zeldovich Approximation to Burgers' Equation: A Plausible Route to a Cellular Automata Adhesion Universe

Victor Christianto<sup>1\*</sup> & Florentin Smarandache<sup>2</sup>

<sup>1</sup>Malang Institute of Agriculture (IPM), Malang, Indonesia

<sup>2</sup>Dept. Mathematics and Sciences, University of New Mexico, NM, USA

### ABSTRACT

Some years ago, Hidding *et al.* suggest that the emergence of an intricate and pervasive web-like structure of the Universe on Megaparsec scales can be approximated by a well-known equation from fluid mechanics, the Burgers' equation. The solution to this equation can be obtained from a geometrical formalism. The resulting Adhesion formalism provides deep insight into the dynamics and topology of the Cosmic Web. It uncovers a direct connection between the conditions in the very early Universe and the complex spatial patterns that emerged out of these under the influence of gravity. In the present paper, we describe a cellular automaton model of the Burgers' equation, which can be investigated via a fast computer simulation. In the end, this suggests a Cellular Automata Adhesion Model of the Universe.

**Keywords:** Discrete physics, cosmology, large scale structure, universe, numerical methods, cellular automata, Burgers equation, Zeldovich approximation.

## 1. Introduction

The Cosmic Web is the fundamental spatial organization of matter on scales of a few up to a hundred Megaparsecs. Galaxies and intergalactic gas matter exist in a wispy web-like arrangement of dense compact clusters, elongated filaments, and sheet-like walls, amidst large near-empty void regions. The filaments are the transport channels along which matter and galaxies flow into massive high-density clusters located at the nodes of the web. The web-like network is shaped by the tidal force field accompanying the inhomogeneous matter distribution [1].

Structure in the Universe has risen out of tiny primordial (Gaussian) density and velocity perturbations by means of gravitational instability. The large-scale anisotropic force field induces anisotropic gravitational collapse, resulting in the emergence of elongated or flattened matter configurations. The simplest model that describes the emergence of structure and complex patterns in the Universe is the Zeldovich Approximation (ZA) [1]. It is our hope that the new approach of the CA Adhesion model of the Universe can be verified either with lab experiments, computer simulation, or by large-scale astronomy observation data.

---

\* Correspondence: Victor Christianto, Malang Institute of Agriculture (IPM), Malang, Indonesia.  
Email: victorchristianto@gmail.com

## 2. From Zeldovich Approximation to Burgers' Equation to the Cellular Automaton model

In this section, we will outline a route from ZA to Burgers' equation and then to the CA model. The simplest model that describes the emergence of structure and complex patterns in the Universe is the Zeldovich Approximation (ZA). In essence, it describes a ballistic flow, driven by a constant (gravitational) potential. The resulting Eulerian position  $x(t)$  at some cosmic epoch  $t$  is specified by the expression [1]:

$$x(t) = q + D(t)u_o(q), \tag{1}$$

where  $q$  is the initial ‘‘Lagrangian’’ position of a particle,  $D(t)$  the time-dependent structure growth factor and

$$u_o = -\nabla_q \Phi_o \tag{2}$$

its velocity. The nature of this approximation may be appreciated by the corresponding source-free equation of motion,

$$\frac{\partial u}{\partial D} + (u \cdot \nabla_x)u = 0. \tag{3}$$

The use of ZA is ubiquitous in cosmology. One major application is its key role in setting up initial conditions in cosmological N-body simulations. Of importance here is its nonlinear extension in terms of the *Adhesion Model* [1].

The ZA breaks down as soon as self-gravity of the forming structures becomes important. To ‘simulate’ the effects of self-gravity, Gurbatov *et al.* included an artificial viscosity. This results in the Burgers' equation as follows [1]:

$$\frac{\partial u}{\partial D} + (u \cdot \nabla_x)u = \nu \cdot \nabla_x^2 u, \tag{4}$$

a well-known PDE from fluid mechanics. This equation has an exact analytical solution, which in the limit of  $\nu \rightarrow 0$ , the solution is [1]:

$$\phi(x, D) = \max_q \left[ \Phi_o(q) - \frac{(x - q)^2}{2D} \right]. \tag{5}$$

This leads to a geometric interpretation of the Adhesion Model. The solution follows from the evaluation of the convex hull of the velocity potential modified by a quadratic term. We found that the solution can also be found by computing the weighted Voronoi diagram of a mesh

weighted with the velocity potential. For more detailed discussion on the Adhesion Model of the Universe, see for example [4].

Now, let us consider another route to solve Burgers' equation: (a) by numerical computation with *Mathematica*, see [3]; and (b) by virtue of the CA approach. Let us skip route (a) and discuss a lesser known approach of cellular automata.

We start with the Burgers' equation with Gaussian white noise that can be rewritten as follows [2]:

$$\frac{\partial u}{\partial t} + \xi = 2u \frac{\partial u}{\partial x} + \frac{\partial^2 u}{\partial x^2} + \eta. \quad (6)$$

By introducing new variables and straightforward calculations afterwards, we have the automata rule [2]:

$$\begin{aligned} \phi_i^{t+1} = & \phi_{i-1}^t + \max[0, \phi_i^t - A, \phi_i^t + \phi_{i+1}^t - B, \Psi_i^t - \phi_{i-1}^t] \\ & - \max[0, \phi_{i-1}^t - A, \phi_{i-1}^t + \phi_i^t - B, \Phi_i^t + \phi_{i-1}^t] \end{aligned} \quad (7)$$

In other words, in this section we give an outline of a plausible route from ZA to Burgers' equation and then to the CA model, which suggests that it appears possible – at least in theory – to consider a nonlinear cosmology based on the CA Adhesion model.

### 3. Concluding Remarks

The use of ZA is ubiquitous in cosmology. One major application is its key role in setting up initial conditions in cosmological N-body simulations. Of importance here is its nonlinear extension in terms of the *Adhesion Model*. In this paper, we give an outline of a plausible route from ZA to Burgers' equation then to the CA model, which suggests that it appears theoretically possible to consider a nonlinear cosmology based on CA Adhesion model.

This paper is part of our theoretical investigation of plausible nonlinear cosmology models beyond Navier-Stokes-inspired approaches.

It is our hope that the proposed approach can be verified with a more extensive computer simulation and (astronomy) observation data.

*Received October 27, 2017; Accepted January 17, 2018*

## References

- [1] J. Hidding, R. van de Weygaert, G. Vegter, B.J.T. Jones, M. Teillaud. The Sticky Geometry of the Cosmic Web. SCG'12, June 17–20, 2012, Chapel Hill, North Carolina, USA. ACM 978-1-4503-1299-8/12/06. arXiv: 1205.1669 [astro-ph.CO] (2012); [1a] J. Hidding, S. Shandarin, R. van de Weygaert. The Zeldovich Approximation: key to understanding Cosmic Web complexity. *Mon. Not. Royal Astron. Soc.* 1-37 (2013)
- [2] Xin-She Yang & Y. Young. Cellular Automata, PDEs, and Pattern Formation. arXiv: 1003.1983 (2010)
- [3] Richard H. Enns & George C. McGuire. *Nonlinear Physics with Mathematica for Scientists and Engineers*. Boston: Birkhäuser, 2001. See pp. 314-316.
- [4] Johan Hidding. *Adhesion: a sticky way of understanding Large Scale Structure*. 2010. 180 p.
- [5] Oliver Hahn. Collisionless Dynamics and the Cosmic Web, a chapter in R. van de Weygaert, S. Shandarin, E. Saar & J. Einasto, eds. *The Zeldovich Universe, Proceedings IAU Symposium No. 308*, 2014. Also in arXiv: 1412.5197 [astro-ph.CO]