

Exploration

On the Emergence of Spacetime Dimensions from Kolmogorov Entropy

Ervin Goldfain *

Advanced Technology and Sensor Group, Welch Allyn Inc., Skaneateles Falls, NY 13153

Abstract

This short pedagogical report is based on a couple of premises. First, it was recently shown that the long run of non-equilibrium Renormalization Group flows is prone to end up on strange attractors. As a result, multifractals are likely to provide the proper framework for the characterization of effective field theories. Secondly, it is known that multifractal analysis uses the Kolmogorov entropy (K-entropy) to quantify the degree of disorder in chaotic systems and turbulent flows. Building on these premises, the report details the remarkable connection between K-entropy, multifractal sets and spacetime dimensions. It also supports the proposal that near and above the Fermi scale, spacetime is defined by continuous and arbitrarily small deviations from four-dimensions.

Keywords: Kolmogorov entropy, multifractals, minimal fractal manifold, effective field theory.

We have conjectured in [1, 2] that the flow from the ultraviolet (UV) to the infrared (IR) sector of any multidimensional nonlinear field theory approaches chaotic dynamics in a universal way. This result stems from several independent routes to aperiodic behavior and implies that the IR attractor of effective field theories is likely to replicate the properties of *strange attractors* (SA) and *multifractal sets*. In particular, the chaotic behavior of the Renormalization Group flow near or above the Fermi scale suggests that phenomena on or above this scale mimic the dynamics on a strange attractor [3-5, 9]. A counterintuitive property of these attractors is that adjacent trajectories separate exponentially fast, yet they remain forever trapped to a bounded region of phase-space. This is to say that, while microscopic SA dynamics is locally unstable, it becomes stable at the global scale. Elaborating further, it can be shown that the global stability of SA trajectories provides the foundation for classical statistical mechanics, which is based on invariant measures and the ergodic hypothesis [10].

Let a generic UV to IR trajectory be described by the n -dimensional phase-space flow $x(\tau)$. Here, τ denotes the evolution parameter (“time”) corresponding to the Renormalization Group scale μ

*Correspondence: Ervin Goldfain, Ph.D., Photonics CoE, Welch Allyn Inc., Skaneateles Falls, NY 13153, USA

E-mail: ervingoldfain@gmail.com

$$\tau = \log\left(\frac{\mu}{\mu_0}\right) \quad (1)$$

The random behavior of the flow near the strange attractor can be characterized by dividing the phase-space into n - dimensional hypercubes of side r , which are sampled at discrete time intervals $\Delta\tau$. The generalized K -entropy of order $q \neq 1$ is given by the equation [6]

$$K_q(X) = -\lim_{r \rightarrow 0} \lim_{\Delta\tau \rightarrow 0} \lim_{N \rightarrow \infty} \frac{1}{N\Delta\tau} \frac{1}{q-1} \ln \sum_{i_1, i_2, \dots, i_N}^{M(r)} p_{i_1, i_2, \dots, i_N}^q \quad (2)$$

where $X = x_i$ is the discrete random variable, that is, $x_i = x(\tau = i\Delta\tau)$, and p_{i_1, i_2, \dots, i_M} stands for the joint probability that the trajectory $x(\tau = \Delta\tau)$ is in i_1 , $x(\tau = 2\Delta\tau)$ is in i_2 and $x(\tau = M\Delta\tau)$ is in i_M . The K -entropy defines the asymptotic scenario where $r \rightarrow 0$ and the phase-space is sampled with an infinite number of steps ($N \rightarrow \infty$) at vanishing time intervals ($\Delta\tau \rightarrow 0$). In the special case $N\Delta\tau = 1$ and when the joint probability is constant across all hypercubes ($M(r) = const.$, $p_{i_1, i_2, \dots, i_M} = p_i$), (2) turns into the *Rényi entropy* in the logarithm base b , which assumes the form

$$S_q(X) = \frac{1}{1-q} \log_b\left(\sum_{i=1}^M p_i^q\right) \quad (3)$$

Furthermore, (3) reduces to the familiar *thermodynamic entropy* when $q \rightarrow 1$ and Boltzmann constant is set to $k_B = 1$ [7]

$$S(X) = -\sum_{i=1}^M p_i \ln p_i \quad (4)$$

Finally, the concept of *generalized dimension* of order q is introduced in conjunction with (3) according to

$$D_q = \lim_{r \rightarrow 0} \frac{1}{1-q} \frac{\log_b\left(\sum_{i=1}^M p_i^q\right)}{\log r} \quad (5)$$

A particularly straightforward expression of (3) is obtained for the null order $q = 0$ and the natural logarithm base. It is referred to as *topological entropy* and is given by

$$S_0(r) = \ln \sum_{i=1}^M p_i^0 = \ln M \quad (6)$$

It is known that the *box-counting dimension* of a fractal object of normalized size r is defined as

$$D_0 \approx \frac{\ln M}{\ln r} \Rightarrow M \approx r^{D_0} = \varepsilon^{-D_0} \quad (7)$$

in which M denotes the number of covering boxes and $\varepsilon = r^{-1}$ is the normalized size of the box. The dimension of ordinary Euclidean space corresponds to integer and positive-definite values of the box-counting dimension, $D_0 = k$, $k = 0, 1, 2, \dots$.

Comparing (6) to (7) leads to the connection between the box-counting dimension and topological entropy via

$$\boxed{\varepsilon^{-D_0} = \exp[S_0(r)]} \quad (8)$$

Two straightforward conclusions may be drawn from (8):

- Maximal topological entropy ($S_0(r) \rightarrow \infty$) matches the limit $\varepsilon \rightarrow 0$ and corresponds to the four-dimensional continuum of both General Relativity and Quantum Field Theory.
- The steady increase of topological entropy along the flow implies that, near or above the Fermi scale, spacetime exhibits a *continuous* spectrum of dimensions $\varepsilon = 4 - D_0 \ll 1$, asymptotically reaching $D_0 = 4$ as $\varepsilon \rightarrow 0$ [8, 4].

Received September 19, 2019; Accepted October 6, 2019

References

- [1] <https://www.prespacetime.com/index.php/pst/article/view/1244> A copy of this article can be found at: https://www.academia.edu/38764569/Multifractal_Analysis_and_the_Dynamics_of_Effective_Field_Theories
- [2] Available at the following site (*in progress*): https://www.academia.edu/38852586/The_Strange_Attractor_Structure_of_Turbulence_and_Effective_Field_Theories_fourth_draft
- [3] Available at the following site: https://www.academia.edu/38735370/Chaotic_Dynamics_of_the_Renormalization_Group_Flow_and_Standard_Model_Parameters
- [4] Available at the following site (*in progress*): https://www.academia.edu/38744111/Bifurcations_and_the_Dynamic_Content_of_Particle_Physics
- [5] <https://arxiv.org/pdf/hep-th/0304178.pdf>
- [6] <https://www.sciencedirect.com/science/article/pii/S0898122113000345>
- [7] <https://arxiv.org/pdf/cond-mat/0207707.pdf>
- [8] <http://www.aracneeditrice.it/index.php/publicazione.html?item=9788854889972>
- Also available at the following site: https://www.researchgate.net/publication/278849474_Introduction_to_Fractional_Field_Theory_consolidated_version

- [9] Available at the following site:

https://www.academia.edu/38752837/Renormalization_group_and_the_emergence_of_random_fractal_topology_in_quantum_field_theory

- [10] Lesne A. and Lagües M, “*Scale Invariance, From Phase Transitions to Turbulence*”, Springer, 2003, p. 325.