

# From Principles to Diagrams in TGD Framework

Matti Pitkänen <sup>1</sup>

## Abstract

In this article, I explore a more explicit realization of twistorialization as lifting of the preferred extremal  $X^4$  of Kähler action to corresponding 6-D twistor space  $X^6$  identified as surface in the 12-D product of twistor spaces of  $M^4$  and  $CP_2$ . Contrary to original expectations, the twistorial approach is not mere reformulation but leads to identification of cosmological constant and perhaps also of gravitational constant based on a first principle and to a modification of the dynamics of Kähler action but preserving the known extremals and basic properties of Kähler action and allowing to interpret induced Kähler form in terms of preferred imaginary unit defining twistor structure. Another new aspect is the fusion of twistorial approach with the vision that diagrams are representations for computations. Also, quantum criticality demands that the diagrams should allow huge symmetries transforming them to braided generalizations of tree-diagrams. Several guiding principles are involved and what is new is the observation that they indeed seem to form a coherent whole.

## 1 Introduction

The generalization of twistor diagrams to TGD framework has been very inspiring and allowed to gain deep insights about what TGD diagrams could be mathematically. Of course, I cannot provide explicit formulas but the general structure for the construction of twistorial amplitudes in  $\mathcal{N} = 4$  SUSY suggests an analogous construction in TGD thanks to huge symmetries of TGD and unique twistorial properties of  $M^4 \times CP_2$ . The twistor program in TGD framework has been summarized in [6].

Contrary to the original expectations, the twistorial approach is not a mere reformulation but leads to a first principle identification of cosmological constant and perhaps also of gravitational constant and to a modification of the dynamics of Kähler action however preserving the known extremals and basic properties of Kähler action and allowing to interpret induced Kähler form in terms of preferred imaginary unit defining twistor structure.

There are some new results forcing a profound modification of the recent view about TGD but consistent with the general picture. A more explicit realization of twistorialization as lifting of the preferred extremal  $X^4$  of Kähler action to corresponding 6-D twistor space  $X^6$  identified as surface in the 12-D product of twistor spaces of  $M^4$  and  $CP_2$  allowing Kähler structure suggests itself. The fiber  $F$  of Minkowskian twistor space must be identified with sphere  $S^2$  with signature  $(-1, -1)$  and would be a variant of the complex space with complex coordinates associated with  $S^2$  and transversal space  $E^2$  in the decomposition  $M^4 = M^2 \times E^2$  and one hyper-complex coordinate associated with  $M^2$ .

The action principle in 6-D context is also Kähler action, which dimensionally reduces to Kähler action plus cosmological term. This brings in the radii of spheres  $S^2(M^4)$  and  $S^2(CP_2)$  associated with the twistor space of  $M^4$  and  $CP_2$ . For  $S(CP_2)$  the radius is of order  $CP_2$  radius  $R$ .  $R(S^2(M^4))$  could be of the order of Planck length  $l_P$ , which would thus become purely classical parameter contrary the expectations. An alternative option is  $R(S^2(M^4)) = R$ . The radius of  $S^2$  associated with space-time surface is determined by the induced metric and is emergent length scale. The normalization of 6-D Kähler action by a scale factor  $1/L^2$  with dimension, which is inverse length squared brings in a further length scale closely related to cosmological constant which is also dynamical and has correct sign to explain accelerated expansion of the Universe. The order of magnitude for  $L$  must be radius of the  $S^2(X^4)$  and therefore small. This could mean a gigantic cosmological constant. Just as in GRT based cosmology!

<sup>1</sup>Correspondence: Matti Pitkänen <http://tgdtheory.fi/>. Address: Karkinkatu 3 | 3, 03600, Karkkila, Finland. Email: [matpitka6@gmail.com](mailto:matpitka6@gmail.com).

This issue can be solved by using the observation that thanks to the decomposition  $H = M^4 \times CP_2$  6-D Kähler action is a sum of two independent terms. The first term corresponds to the 6-D lift of the ordinary Kähler action and for it the contribution from  $S^2(CP_2)$  fiber is assumed to be absent: this could be due to the imbedding of  $S^2(X^4)$  reducing to identification  $S^2(M^4)$  and is not true generally. Second term in action is assumed to come from the  $S^2(M^4)$  fiber of twistor space  $T(M^4)$ . The independency implies that couplings strengths are independent for them.

The analog for Kähler coupling strength (analogous to critical temperature) associated with  $S^2(M^4)$  must be extremely large - so large that one has  $\alpha_K(M^4) \times R(M^4)^2 \sim L^2$ ,  $L$  size scale of the recent Universe. This makes possible the small value of cosmological constant assignable to the volume term given by this part of the dimensionally reduced action. Both Kähler coupling strengths are assumed to have a spectrum determined by quantum criticality and the spectrum of  $\alpha_K(M^4)$  comes essentially as p-adic primes satisfying p-adic length scale hypothesis  $p \simeq 2^k$ ,  $k$  prime. In fact, it turns that one can assume that the entire 6-D Kähler action contributes if one assumes that the winding numbers  $(w_1, w_2)$  for the map  $S^2(X^4) \rightarrow S^2(M^4) \times S^2(CP_2)$  satisfy  $(w_1, w_2) = (n, 0)$  in cosmological scales. The identification of  $w_1$  as  $h_{eff}/h = n$  is highly suggestive.

The dimensionally reduced dynamics is a highly non-trivial modification of the dynamics of Kähler action however preserving the known extremals and basic properties of Kähler action and allowing to interpret induced Kähler form in terms of preferred imaginary unit defining twistor structure. Strong constraints come also from the condition that induced spinor structure coming from that for twistor space  $T(H)$  is essentially that coming from that of  $H$ .

Second new element is the fusion of the twistorial approach with the vision that diagrams are representations for computations. This as also quantum criticality demands that the diagrams should allow huge symmetries allowing to transform them to braided generalizations of tree-diagrams. Several guiding principles are involved and what is new is the observation that they indeed seem to form a coherent whole.

In the sequel I will discuss the recent understanding of twistorialization, which is considerably improved from that in the earlier formulation. I formulate the dimensional reduction of 6-D Kähler action and consider the physical interpretation. There are considerable uncertainties at the level of details I dare believe that basically the situation is understood. After that I proceed to discuss the basic principles behind the recent view about scattering amplitudes as generalized Feynman diagrams.

## 2 Twistorial lift of Kähler action

First I will try to clarify the mathematical details related to the twistor spaces and how they emerge in the recent context. I do not regard myself as a mathematician in technical sense and I can only hope that the representation based on physical intuition does not contain serious mistakes.

### 2.1 Imbedding space is twistorially unique

It took roughly 36 years to learn that  $M^4$  and  $CP_2$  are twistorially unique. Space-times are surfaces in  $H = M^4 \times CP_2$ .  $M^4$  and  $CP_2$  are unique 4-manifolds in the sense that both allow twistor space with Kähler structure: Kähler structure is the crucial concept. Strictly speaking, it is  $E^4$  and  $S^4$  allow twistor space with Kähler structure: in the case of  $M^4$  signature could cause problems. The standard identification for the twistor space of  $M^4$  would be Minkowskian variant  $PT = P_3 = SU(2, 2)/SU(2, 1) \times U(1)$  of 6-D twistor space  $PT = CP_3 = SU(4)/SU(3) \times U(1)$  of  $E^4$ . The twistor space of  $CP_2$  is 6-D  $T(CP_2) = SU(3)/U(1) \times U(1)$ , the space for the choices of quantization axes of color hypercharge and isospin.

The case of  $M^4$  is however problematic. It is often stated that the twistor space is  $PT = CP_3 = SU(4)/SU(3) \times U(1)$ . The metric of twistor space does not appear in the construction of twistor amplitudes. Already the basic structure of  $PT$  suggests that this identification cannot be correct.

As if the situation were not complicated enough, there are two notions of twistor space: the twistor space identified as  $P_3$  and as a trivial sphere bundle  $M^4 \times CP_1$  having Kähler structure - what Kähler

structure actually means in case of  $M^4$  is however not quite clear.

These considerations lead to a proposal - just a proposal - for the formulation of TGD in which space-time surfaces  $X^4$  in  $H$  are lifted to twistor spaces  $X^6$ , which are sphere bundles over  $X^4$  and such that they are surfaces in 12-D product space  $T(M^4) \times T(CP_2)$  such the twistor structure of  $X^4$  are in some sense induced from that of  $T(M^4) \times T(CP_2)$ . In the following  $T(M^4)$  therefore denotes the trivial sphere bundle  $M^4 \times CP_1$  over  $M^4$  and twistorialization of scattering amplitudes would involve the projection from  $T(M^4)$  to  $P_3$ . What is nice in this formulation is that one couldn use all the machinery of algebraic geometry so powerful in superstring theory (Calabi-Yau manifolds).

## 2.2 Some basic definitions

What twistor structure in Minkowskian signature does really mean geometrically has remained a confusing question for me. The problems associated with the Minkowskian signature of the metric are encountered also in twistor Grassmann approach to the scattering amplitudes but are circumvented by performing Wick rotation that is using  $E^4$  or  $S^4$  instead of  $M^4$  and applying algebraic continuation. Also complexification of Minkowski space for momenta is used. These tricks do not apply now.

To make this more concrete, let us sum up the basic definitions.

1. Bi-spinors in representations  $(1/2,0)$  and  $(0,1/2)$  of Lorentz group are the building bricks of twistors. Bi-spinors  $v^a$  and their conjugates  $v^{a'}$  have the following inner products:

$$\begin{aligned} \langle vw \rangle &= \epsilon_{ab} v^a w^b, & [vw] &= \epsilon_{a'b'} v^{a'} w^{b'}, \\ \epsilon_{ab} &= (0, 1; -1, 0), & \epsilon_{a'b'} &= (0, 1; -1, 0). \end{aligned} \quad (2.1)$$

Unprimed spinor and its primed variant of the spinor are related by complex conjugation. Index raising is by the inverse  $\epsilon^{ab}$  of  $\epsilon_{ab}$ .

2. Twistors are identified as pairs of 2-spinor and its conjugate

$$Z^\alpha = (\lambda_a, \mu^{a'}), \quad \bar{Z}_\alpha = (\bar{\mu}^a, \lambda_{a'}) \quad (2.2)$$

The norm for  $Z^\alpha$  is defined as

$$Z^\alpha \bar{Z}^\alpha = \langle \lambda \bar{\mu} \rangle + [\bar{\lambda} \mu]. \quad (2.3)$$

One can write the metric explicitly as direct sum of terms of form  $dudv$  (metric of  $M^2$ ) and each of the can be taken to diagonal form  $(1,-1)$ . Hence the metric can be written as  $diag(1, 1, 1, 1, -1, -1, -1, -1)$ .

3. This norm allows to decompose  $PT$  to 3 parts  $PT_+, PT_-$  and  $PN$  in a projectively invariant manner depending on whether the sign of the norm is negative, positive, or whether it vanishes.  $PT_+$  and  $PT_-$  serve as loci for the twistorial lifts of positive and negative energy modes of massless fields.  $PN$  corresponds to the 5-D boundary of the lightcone of  $M(2, 4)$ . By projective identification along light-like radial coordinate it reduces to what is known as conformal compactification of  $M^4$ , whose metric is defined only apart from a conformal factor. The natural metric of  $PT = P_3$  does not seem to play any role in the construction of the amplitudes relying on projective invariants. The signature of  $M^4$  metric however makes itself visible in the structure of  $PT$ : for the Euclidian variant of twistor space one would not have this decomposition to three parts.

Another definition of twistor space - to be used in the geometrization of twistor approach to be proposed - is as a trivial  $S^2$  bundle  $M^4 \times CP_1$  over  $M^4$ . Since the twistor spheres associated with the points of  $M^4$  with light-like separation intersect, these two definitions cannot be equivalent. In fact, the proper definition of twistor space relies on double fibration involving both views about twistor space discussed in [2] (see <http://arxiv.org/pdf/1308.2820.pdf>).

1. The twistor bundle denoted as  $PS$  is the product  $M^4 \times CP_1$  with  $CP_1$  realized as projective space and having coordinates  $(x^{aa'}, \lambda_a)$ ,  $\{x^{aa'}\} \leftrightarrow x^\mu \sigma_\mu$ , where the spinor  $\lambda_a$  is projective 2-spinor in  $(1/2, 0)$  representation.
2. The twistors defined in this manner have a trivial projection  $q$  to  $M^4$  and non-trivial projection  $p$  to  $P_3$  with local projective coordinates  $(\lambda_a, \mu^{a'})$ . The projection  $p$  is defined by the projectively invariant incidence relation

$$\mu^{a'} = ix^{aa'} \lambda_a$$

If  $y^{aa'}$  and  $a^{aa'}$  differ by light-like vector there exists spinor  $\lambda$  annihilated by the the difference vector and there exists twistor  $(\lambda_a, \mu^{a'})$  to which both  $(x, \lambda)$  and  $(y, \lambda)$  are mapped by the incidence relation. Thus the images of twistor spheres associated for points with light-like separation intersect so that one does not have a proper  $CP_1$  bundle structure.

3. The trivial twistor bundle  $T(M^4) = M^4 \times CP_1$  would define the twistor space of  $M^4$  in geometric sense. For this space the metric matters and the radius of  $CP_1$  turns out to allow identification in terms of Planck length. Gravitational interaction would bring in Planck length as a basic scale in this manner.  $PT$  in turn would define the twistor space in which the twistorial lifts of imbedding space-spinor fields are defined. For this space the metric, which is degenerate and seems to be only projectively defined should not be relevant as the construction of twistorial amplitudes suggests. Note however that the identification as the Minkowskian variant of  $P_3$  allows also the introduction of metric.

This picture has an important immediate implication for the construction of quantum TGD. Positive and negative energy parts of zero energy states are defined at light-like boundaries of  $CD \times CP_2$ , where  $CD$  is the intersection of future and past directed light-cones. The twistorial lifts of the amplitudes from  $\delta CD \times CP_2$  must be single valued. The strongest condition guaranteeing this is that they do not depend on the radial light-like coordinate at  $\delta CD$ . Super-symplectic symmetry implying the analog of conformal gauge symmetry for the radial light-like coordinate could guarantee this. There is however a hierarchy of conformal gauge symmetry breakings corresponding to the inclusion hierarchy of isomorphic sub-algebras so that this condition is too strong. A weaker condition is that the amplitude  $F(m, \lambda)$  in  $T(M^4)$  is constant along the light-like ray for the  $\lambda$  associated with the  $m$  along this ray. An even stronger condition is that  $F(m, \lambda)$  vanishes along the ray. Particle would not propagate along  $\delta CD$  and would avoid remaining at the boundary of  $CD$ , a condition which is perfectly sensible physically.

### 2.3 What does twistor structure in Minkowskian signature really mean?

The following considerations relate to  $T(M^4)$  identified as trivial bundle  $M^4 \times CP_1$  with natural coordinates  $(m^{aa'}, \lambda_a)$ , where  $\lambda_a$  is projective spinor. The challenge is to generalize the complex structure of twistor space of  $E^4$  to that for  $M^4$ . It turns out that the assumption that twistor space has ordinary complex structure fails. The first guess was that the fiber of twistor space is hyperbolic sphere with metric signature  $(1, -1)$  having infinite area so that the 6-D Kähler action would be infinite. This makes no sense. The only alternative, which comes in mind is a hypercomplex generalization of the Kähler structure for  $M^4$  lifted to twistor space, which locally means only adding of  $S^2$  fiber with metric signature  $(-1, -1)$ .

1. To proceed one must make an explicit the definition of twistor space. The 2-D fiber  $S^2$  consists of antisymmetric tensors of  $X^4$  which can be taken to be self-dual or anti-self-dual by taking any antisymmetric form and by adding to its plus/minus its dual. Each tensor of this kind defines a direction - point of  $S^2$ . These points can be also regarded as quaternionic imaginary units. One has a natural metric in  $S^2$  defined by the  $X^4$  inner product for antisymmetric tensors: this inner product depends on space-time metric. Kähler action density is example of a norm defined by this inner product in the special case that the antisymmetric tensor is induced Kähler form. Induced Kähler form defines a preferred imaginary unit and is needed to define the imaginary part  $\omega(X, Y) = ig(X, -JY)$  of hermitian form  $h = h + i\omega$ .
2. To define the analog of Kähler structure for  $M^4$ , one must start from a decomposition of  $M^4 = M^2 \times E^2$  ( $M^2$  is generated by light-like vector and its dual) and  $E^2$  is orthogonal to it.  $M^2$  allows hypercomplex structure, which light-like coordinates ( $u = t - z, v = t + z$ ) and  $E^2$  complex structure and the metric has form  $ds^2 = dudv + dzd\bar{z}$ . Hypercomplex numbers can be represented as  $h = t + iez, i^2 = -1, e^2 = -1, i^2 = -1, e^2 = -1$ . Hyper-complex numbers do not define number field since for light-like hypercomplex numbers  $t + iez, t = \pm z$  do not have finite inverse. Hypercomplex numbers allow a generalization of analytic functions used routinely in physics. Kähler form representing hypercomplex imaginary unit would be replaced with  $eJ$ . One would consider subspaces of complexified quaternions spanned by real unit and units  $eI_k, k = 1, 2, 3$  as representation of the tangent space of space-time surfaces in Minkowskian regions. This is familiar already from  $M^8$  duality [8].

$M^4 = M^2 \times E^2$  decomposition can depend on point of  $M^4$  (polarization plane and light-like momentum direction depend on point of  $M^4$ ). The condition that this structure allows global coordinates analogous to  $(u, v, z, \bar{z})$  requires that the distributions for  $M^2$  and  $E^2$  are integrable and thus define 2-D surfaces. I have christened this structure Hamilton-Jacobi structure. It emerges naturally in the construction of extremals of Kähler action that I have christened massless extremals (MEs,[3]) and also in the proposal for the generalization of complex structure to Minkowskian signature [9].

One can define the analog of Kähler form by taking sum of induced Kähler form  $J$  and its dual  $*J$  defined in terms of permutation tensor. The normalization condition is that this form integrates to the negative of metric  $(J \pm *J)^2 = -g$ . This condition is possible to satisfy.

3. How to lift the Hamilton Jacobi structure of  $M^4$  to Kähler structure of its twistor space? The basic definition of twistors assumes that their exists a field of time-like directions, and that one considers projections of 4-D antisymmetric tensors to the 3-space orthogonal to the time-like direction at given point. One can say that the projection yields magnetic part of the antisymmetric tensor (say induced Kähler form  $J$ ) with positive norm with respect to natural metric induced to the twistor fiber from the inner product between two-forms. This unique time direction would be defined the light-like vector defining  $M^2$  and its dual. Therefore the signature of the metric of  $S^2$  would be  $(-1, -1)$ . In quaternionic picture this direction corresponds to real quaternionic unit.
4. To sum up, the metric of the Minkowskian twistor space has signature  $(-1, -1, 1, -1, -1, -1)$ . The Minkowskian variant of the twistor space would give 2 complex coordinates and one hypercomplex coordinate. Cosmological term would be finite and the sign of the cosmological term in the dimensionally reduced action would be positive as required. Also metric determinant would be imaginary as required. At this moment I cannot invent any killer objection against this option.

It must be made clear that the proposed definition of twistor space of  $M^4$  does not seem to be equivalent with the twistor space assignable to conformally compactified  $M^4$ . One has trivial  $S^2$  bundle and Hamilton-Jacobi structure, which is hybrid of complex and hyper-complex structure.

## 2.4 What does the induction of the twistor structure to space-time surface really mean?

Consider now what the induction of the twistor structure to space-time surface  $X^4$  could mean.

1. The induction procedure for Kähler structure of 12-D twistor space  $T$  requires that the induced metric and Kähler form of the base space  $X^4$  of  $X^6$  obtained from  $T$  is the same as that obtained by inducing from  $H = M^4 \times CP_2$ . Since the Kähler structure and metric of  $T$  is lift from  $H$  this seems obvious. Projection would compensate the lift.
2. This is not yet enough. The Kähler structure and metric of  $S^2$  projected from  $T$  must be same as those lifted from  $X^4$ . The connection between metric and  $\omega$  implies that this condition for Kähler form is enough. The antisymmetric Kähler forms in fiber obtained in these two manners co-incide. Since Kähler form has only one component in 2-D case, one obtains single constraint condition giving a commutative diagram stating that the direct projection to  $S^2$  equals with the projection to the base followed by a lift to fiber. The resulting induced Kähler form is not covariantly constant but in fiber  $S^2$  one has  $J^2 = -g$ .

As a matter of fact, this condition might be trivially satisfied as a consequence of the bundle structure of twistor space. The Kähler form from  $S^2 \times S^2$  can be projected to  $S^2$  associated with  $X^4$  and by bundle projection to a two-form in  $X^4$ . The intuitive guess - which might be of course wrong - is that this 2-form must be same as that obtained by projecting the Kähler form of  $CP_2$  to  $X^4$ . If so then the bundle structure would be essential but what does it really mean?

3. Intuitively it seems clear that  $X^6$  must decompose locally to a product  $X^4 \times S^2$  in some sense. This is true if the metric and Kähler form reduce to direct sums of contributions from the tangent spaces of  $X^4$  and  $S^2$ . This guarantees that 6-D Kähler action decomposes to a sum of 4-D Kähler action and Kähler action for  $S^2$ .

This could be however too strong a condition. Dimensional reduction occurs in Kaluza-Klein theories and in this case the metric can have also components between tangent spaces of the fiber and base being interpreted as gauge potentials. This suggests that one should formulate the condition in terms of the matrix  $T \leftrightarrow g^{\alpha\mu}g^{\beta\nu} - g^{\alpha\nu}g^{\beta\mu}$  defining the norm of the induced Kähler form giving rise to Kähler action.  $T$  maps Kähler form  $J \leftrightarrow J_{\alpha\beta}$  to a contravariant tensor  $J_c \leftrightarrow J^{\alpha\beta}$  and should have the property that  $J_c(X^4) (J_c(S^2))$  does not depend on  $J(S^2) (J(X^4))$ .

One should take into account also the self-duality of the form defining the imaginary unit. In  $X^4$  the form  $S = J \pm *J$  is self-dual/anti-self dual and would define twistorial imaginary unit since its square equals to  $-g$  representing the negative of the real unit. This would suggest that 4-D Kähler action is effectively replaced with  $(J \pm *J) \wedge (J \pm *J) = J^*J \pm J \wedge J$ , where  $*J$  is the Hodge dual defined in terms of 4-D permutation tensor  $\epsilon$ . The second term is topological term (Abelian instanton term) and does not contribute to field equations. This in turn would mean that it is the tensor  $T \pm \epsilon$  for which one can demand that  $S_c(X^4) (S_c(S^2))$  does not depend on  $S(S^2) (S(X^4))$ .

4. The preferred quaternionic imaginary unit should be represented as a projection of Kähler form of 12-D twistor space  $T(H)$ . The preferred imaginary unit defining twistor structure as sum of projections of both  $T(CP_2)$  and  $T(M^4)$  Kähler forms would guarantee that vacuum extremals like canonically imbedded  $M^4$  for which  $T(CP_2)$  Kähler form contributes nothing have well-defined twistor structure.  $T(M^4)$  or  $T(CP_2)$  are treated completely symmetrically but the maps of  $S^2(X^4)$  to  $S^2(M^4)$  and  $S^2(CP_2)$  characterized by winding numbers induce symmetry breaking.

For Kähler action  $M^4 - CP_2$  symmetry does not make sense. 4-D Kähler action to which 6-D Kähler action dimensionally reduces can depend on  $CP_2$  Kähler form only. I have also considered the possibility of covariantly constant self-dual  $M^4$  term in Kähler action but given it up because of problems with Lorentz invariance. One should couple the gauge potential of  $M^4$  Kähler form to

induced spinors. This would mean the existence of vacuum gauge fields coupling to sigma matrices of  $M^4$  so that the gauge group would be non-compact  $SO(3, 1)$  leading to a breakdown of unitarity.

There is still one difficulty to be solved.

1. The normalization of 6-D Kähler action by a scale factor  $1/L^2$  with dimension, which is inverse length squared, brings in a further length scale. The first guess is that  $1/L^2$  is closely related to cosmological constant, which is also dynamical and  $1/L^2$  has indeed correct sign to explain accelerated expansion of the Universe. Unfortunately, if  $1/L^2$  is of order cosmological constant, the value of the ordinary Kähler coupling strength  $\alpha_K$  would be enormous. As a matter of fact, the order of magnitude for  $L^2$  must be equal to the area of  $S^2(X^4)$  and in good approximation equal to  $L^2 = 4\pi R^2(S^2(M^4))$  and therefore in the same range as Planck length  $l_P$  and  $CP_2$  radius  $R$ . This would imply a gigantic value of cosmological constant. Just as in GRT based cosmology!
2. This issue can be solved by using the observation that thanks to the decomposition  $H = M^4 \times CP_2$ , 6-D Kähler action is sum of two independent terms. The first term corresponds to the 6-D lift of the ordinary Kähler action. For it the contribution from  $S^2(CP_2)$  fiber is absent if the imbedding of  $S^2(X^4)$  to  $S^2(M^4) \times S^2(CP_2)$  reduces to identification with  $S^2(M^4)$  so that  $S^2(CP_2)$  is effectively absent: this is not true generally. Second term in the action is assumed to come from the  $S^2(M^4)$  fiber of twistor space  $T(M^4)$ , which can indeed contribute without breaking of Lorentz symmetry. In fact, one can assume that also the Kähler form of  $M^4$  contributes as will be found.
3. The independency implies that Kähler couplings strengths are independent for them. If one wants that cosmological constant has a reasonable order of magnitude,  $L \sim R(S^2(M^4))$  must hold true and the analog  $\alpha_K(S^2(M^4))$  of the ordinary Kähler coupling strength (analogous to critical temperature) must be extremely large - so large that one has

$$\alpha_K(M^4) \times 4\pi R(M^4)^2 \sim L^2 \quad ,$$

where  $L$  is the size scale of the recent Universe.

This makes possible the small value of cosmological constant assignable to the volume term given by this part of dimensionally reduced action. Both Kähler coupling strengths are assumed to have a spectrum determined by quantum criticality and the spectrum of  $\alpha_K(M^4)$  would be essentially as p-adic primes satisfying p-adic length scale hypothesis  $p \simeq 2^k$ ,  $k$  prime. One can criticize this identification of 6-D Kähler action as artificial but it seems to be the only option that works. Interestingly also the contribution from  $M^4$  Kähler form can be allowed since it is also extremely small. For canonically imbedded  $M^4$  this contribution vanishes by self-duality of  $M^4$  Kähler form and is extremely small for the vacuum extremals of Kähler action.

4. For general winding numbers of the map  $S^2(X^4) \rightarrow S^2(M^4) \times S^2(CP_2)$  also  $S^2(CP_2)$  Kähler form contributes and cosmological constant is gigantic. It would seem that only the winding numbers  $(w_1, w_2) = (n, 0)$  are consistent with the observed value of cosmological constant. Hence it seems that there is no need to pose any additional conditions to the Kähler action if one uses the fact that  $T(M^4)$  and  $T(CP_2)$  parts are independent!

It is good to list the possible open issues related to the precise definition of the twistor structure and of  $M^4$  Kähler action.

1. The proposed definition of  $M^4$  twistor space a Cartesian product of  $M^4$  and  $S^2(M^4)$  parts involving Hamilton-Jacobi structure does not seem to be equivalent with the twistor identification as  $SU(2, 2)/SU(2, 1) \times U(1)$  having conformally compactified  $M^4$  as base space. There exists an entire moduli space of Hamilton-Jacobi structures. If the  $M^4$  part of Kähler form participates in dynamics, one must include the specification of the Hamilton-Jacobi structure to the definition of CD and

integrate over Hamilton Jacobi-structures as part of integral over WCW in order to gain Lorentz invariance. Note that Hamilton-Jacobi structure enters to dynamics also through the construction of massless extremals [3].

2. The presence of  $M^4$  part of Kähler form in action implies breaking of Lorentz invariance for extremals of lifted Kähler action. The same happens at the level of induced spinors if this Kähler form couples to imbedding space spinors. If  $T(M^4)$  is trivial bundle, one can include only the  $T(S^2(M^4))$  part of Kähler form to Kähler action and couple only this to the spinors of  $T(H)$ . The integration over Hamilton-Jacobi structures becomes un-necessary.
3. If one includes  $M^4$  part of Kähler form to 6-D Kähler action, one has several options. One can have sum of the Kähler actions for  $T(M^4)$  and  $T(CP_2)$  or Kähler action defined by the sum  $J(T(M^4))/g_K$  and  $J(T(CP_2))/\alpha_K$  with  $\alpha_K(M^4) = g_K^2(M^4)/4\pi\hbar$  and  $\alpha_K = g_K^2/4\pi\hbar$  with a proper normalization to guarantee that the squares of induced Kähler forms give sum of Kähler actions as in the first option. In this case one obtains interference term proportional to  $Tr(J(M^4)J(CP_2))$ . For the proposed value of  $\alpha_K$  also the interference term is extremely small as compared to Kähler action in recent cosmology.

## 2.5 Could $M^4$ Kähler form introduce new gravitational physics?

The introduction of  $M^4$  Kähler form could bring in new gravitational physics.

1. As found, the twistorial formulation of TGD assigns to  $M^4$  a self dual Kähler form whose square gives Minkowski metric. It can (but need not if  $M^4$  twistor space is trivial as bundle) contribute to the 6-D twistor counterpart of Kähler action inducing  $M^4$  term to 4-D Kähler action vanishing for canonically imbedded  $M^4$ .
2. Self-dual Kähler form in empty Minkowski space satisfies automatically Maxwell equations and has by Minkowskian signature and self-duality a vanishing action density. Energy momentum tensor is proportional to the metric so that Einstein Maxwell equations are satisfied for a non-vanishing cosmological constant!  $M^4$  indeed allows a large number of self dual Kähler fields (I have christened them as Hamilton-Jacobi structures). These are probably the simplest solutions of Einstein-Maxwell equations that one can imagine!
3. There however exist quite a many Hamilton-Jacobi structures. However, if this structure is to be assigned with a causal diamond (CD) it must satisfy additional conditions, say  $SO(3)$  symmetry and invariance under time translations assignable to CD. Alternatively, covariant constancy and  $SO(2) \subset SO(3)$  symmetry might be required.

This raises several questions. Could  $M^4$  Kähler form replace  $CP_2$  Kähler form in the picture for how gravitational interaction is mediated at quantal level? Could one speak of flux tubes of the magnetic part of this Kähler form? Or should one consider the Kähler field as a sum of the two Kähler forms weighted by the inverses  $1/g_K$  of corresponding Kähler couplings. If so then  $M^4$  contribution would be negligible except for canonically imbedded  $M^4$  in the recent cosmology. Note that  $\alpha_K$  and  $\alpha_K(M^4)$  have interpretation as analogs of quantum critical temperatures but can depend on the p-adic lengths scale defining the cosmology.

1. The natural expectation is that Kähler form characterizes CD having preferred time direction suggested strongly by number theoretical considerations involving quaternionic structure with preferred direction of time axis assignable to real unit quaternion.

Self-duality gives rise to Kähler magnetic and electric fields in the same spatial direction identifiable as a local quantization axis for spin assignable to CD assignable to observer. CD indeed serves as a correlate for conscious entity in TGD inspired theory of consciousness. Flux tube would connect

mass  $M$  to mass  $m$  assignable to observer and flux tube direction would define spin quantization axes for the CD of the observer. Spin quantization axis would be naturally in the direction of magnetic field, which is direction of the flux tube.

2. The self-dual Kähler form could be spherically symmetric for CDs and represent self dual magnetic monopole field (dyon) with monopole charge at the line connecting the tips of CD and have non-vanishing components  $J^{tr} = \epsilon^{tr\theta\phi} J_{\theta\phi}$ ,  $J_{\theta\phi} = \sin(\theta)$ . One would have genuine monopole, which is somewhat questionable feature. Only the entire radial flux would be quantized. CD could be associated with the mass  $M$  of the central object. The gauge potential associated with  $J$  could be chosen to be  $A_\mu \leftrightarrow (1/r, 0, 0, \cos(\theta))$ . I have considered this kind of possibility earlier in context of TGD inspired model of anyons but gave up the idea.

The moduli space for CDs with second tip fixed would be hyperbolic space  $H^3 = SO(3, 1)/SO(3)$  or a space obtained by identifying points at the orbits of some discrete subgroup of  $SO(3, 1)$  as suggested by number theoretic considerations. This induced Kähler field could make the blackholes with center at this line to behave like  $M^4$  magnetic monopoles if the  $M^4$  part of Kähler form is induced into the 6-D lift of Kähler action with extremely small coefficients of order of magnitude of cosmological constant. Cosmological constant and the possibility of CD monopoles would thus relate to each other.

3. The self-dual  $M^4$  Kähler form could be also covariantly constant ( $J_{tz} = J_{xy} = 1$ ) and represent electric and magnetic fluxes in a fixed direction identifiable as a quantization axes for spin and characterizing CD. In this case the CD would be associated with the mass  $m$  of observer. The moduli space of CDs would be now  $SO(3, 1)/SO(1, 1) \times SO(2)$  which is completely analogous to the twistor space  $SU(3)/U(1) \times U(1)$ .
4. Boundary conditions (allowing no boundaries!) demand that the flux tubes have closed cross section - say sphere  $S^2$  - rather than disk: stability is guaranteed if the  $S^2$  cross section is mapped to homologically non-trivial surface of  $CP_2$  or is projection of it. This would give monopole flux also for  $CP_2$  Kähler form so that the original hypothesis would be correct.
5. Radial flux tubes are possible both spherically symmetric and covariantly constant Kähler form possibly mediating gravitational interaction but the flux is not quantized unless preferred extremal property implies this: in any case  $M^4$  flux would be very small unless one has large value of gravitational Planck constant implying  $n$ -sheeted covering of  $M^4$  and flux is scale up by  $n$  since every sheet gives a contribution. For spherically symmetric  $M^4$  Kähler form the flux tubes would have naturally conical structure spanning a constant solid angle. For covariantly constant Kähler form the flux tubes would be cylindrical.

There are further interpretational problems.

1. The classical coupling of  $M^4$  Kähler gauge potential to induced spinors is not small. Can one really tolerate this kind of coupling equivalent to a coupling to a self dual monopole field carrying electric and magnetic charges? One could of course consider the condition that the string world sheets carrying spinor modes are such that the induced  $M^4$  Kähler form vanishes and gauge potential become pure gauge.  $M^4$  projection would be 2-D Lagrange manifold whereas  $CP_2$  projection would carry vanishing induced  $W$  and possibly also  $Z^0$  field in order that em charge is well defined for the modes. These conditions would fix the string world sheets to a very high degree in terms of maps between this kind of 2-D sub-manifolds of  $M^4$  and  $CP_2$ . Spinor dynamics would be determined by the avoidance of interaction!

Recall that one could interpret the localization of spinor modes to 2-surfaces in the sense of strong form of holography: one can continue induced spinor fields to the space-time interior as indeed assumed but the continuation is completely determined by the data at 2-D string world sheets.

It must be emphasized that the imbedding space spinor modes characterizing the ground states of super-symplectic representations would not couple to the monopole field so that at this level Poincare invariance is not broken. The coupling would be only at the space-time level and force spinor modes to Lagrangian sub-manifolds.

2. At the static limit of GRT and for  $g_{ij} \simeq \delta_{ij}$  implying  $SO(3)$  symmetry there is very close analogy with Maxwell's equations and one can speak of gravi-electricity and gravi-magnetism with 4-D vector potential given by the components of  $g_{t\alpha}$ . The genuine  $U(1)$  gauge potential does not however relate to the gravimagnetism in GRT sense. Situation would be analogous to that for  $CP_2$ , where one must add to the spinor connection  $U(1)$  term to obtain respectable spinor structure. Now the  $U(1)$  term would be added to trivial spinor connection of flat  $M^4$ : its presence would be justified by twistor space Kähler structure. If the induced  $M^4$  Kähler form is present as a classical physical field it means genuinely new contribution to  $U(1)$  electroweak of standard model. If string world sheets carry vanishing  $M^4$  Kähler form, this contribution vanishes classically.

## 2.6 A connection with the hierarchy of Planck constants?

A connection with the hierarchy of Planck constants is highly suggestive. Since also a connection with the p-adic length scale hierarchy suggests itself for the hierarchy of p-adic length scales it seems that both length scale hierarchies might find first principle explanation in terms of twistorial lift of Kähler action.

1. Cosmological considerations encourage to think that  $R_1 \simeq l_P$  and  $R_2 \simeq R$  hold true. One would have in early cosmology  $(w_1, w_2) = (1, 0)$  and later  $(w_1, w_2) = (0, 1)$  guaranteeing  $R_D$  grows from  $l_P$  to  $R$  during cosmological evolution. These situations would correspond the solutions  $(w_1 = n, 0)$  and  $(0, w_2 = n)$  one has  $A = n4\pi R_1^2$  and  $A = n \times 4\pi R_2^2$  and both Kähler coupling strengths are scaled down to  $\alpha_K/n$ . For  $\hbar_{eff}/h = n$  exactly the same thing happens!

There are further intriguing similarities.  $\hbar_{eff}/h = n$  is assumed to correspond *multi-sheeted* (to be distinguished from *many-sheeted*!) covering space structure for space-time surface. Now one has covering space defined by the lift  $S^2(X^4) \rightarrow S^2(M^4) \times S^2(CP_2)$ . These lifts define also lifts of space-time surfaces.

Could the hierarchy of Planck constants correspond to the twistorial surfaces for which  $S^2(M^4)$  is  $n$ -fold covering of  $S^2(X^4)$ ? The assumption has been that the  $n$ -fold multi-sheeted coverings of space-time surface for  $\hbar_{eff}/h = n$  are singular at the ends of space-time surfaces at upper and lower boundaries if causal diamond (CD). Could one consider a more precise definition of twistor space in such a manner that  $CD$  replaces  $M^4$  and the covering becomes singular at the light-like boundaries of CD - the branches of space-time surface would collapse to single one.

Does this collapse have a clear geometric meaning? Are the projections of various branches of the  $S^2$  lift automatically identical so that one would have the original picture in which one has  $n$  identical copies of the same space-time surface? Or can one require identical projections only at the light-like boundaries of CD?

2.  $w_1 = w_2 = w$  is essentially the first proposal for conditions associated with the lifting of twistor space structure.  $w_1 = w_2 = n$  gives  $ds^2 = (R_1^2 + R_2^2)(d\theta^2 + w^2 d\phi^2)$  and  $A = n \times 4\pi(R_1^2 + R_2^2)$ . Also now Kähler coupling strength is scaled down to  $\alpha/n$ . Again a connection with the hierarchy of Planck constants suggests itself.
3. One can consider also the option  $R_1 = R_2$  option giving  $ds^2 = R_1^2(2d\theta^2 + (w_1^2 + w_2^2)d\phi^2)$ . If the integers  $w_i$  define Pythagorean square one has  $w_1^2 + w_2^2 = n^2$  and one has  $R_1 = R_2$  option that one has  $A = n \times 4\pi R^2$ . Also now the connection with the hierarchy of Planck constants might make sense.

## 2.7 Twistorial variant for the imbedding space spinor structure

The induction of the spinor structure of imbedding space is in key role in quantum TGD. The question arises whether one should lift also spinor structure to the level of twistor space. If so one must understand how spinors for  $T(M^4)$  and  $T(CP_2)$  are defined and how the induced spinor structure is induced.

1. In the case of  $CP_2$  the definition of spinor structure is rather delicate and one must add to the ordinary spinor connection  $U(1)$  part, which corresponds physically to the addition of classical  $U(1)$  gauge potential and indeed produces correct electroweak couplings to quarks and leptons. It is assumed that the situation does not change in any essential manner: that is the projections of gauge potentials of spinor connection to the space-time surface give those induced from  $M^4 \times CP_2$  spinor connection plus possible other parts coming as a projection from the fiber  $S^2(M^2) \times S^2(CP_2)$ . As a matter of fact, these other parts should vanish if dimensional reduction is what it is meant to be.
2. The key question is whether the complications due to the fact that the geometries of twistor spaces  $T(M^4)$  and  $T(CP_2)$  are not quite Cartesian products (in the sense that metric could be reduced to a direct sum of metrics for the base and fiber) can be neglected so that one can treat the sphere bundles approximately as Cartesian products  $M^4 \times S^2$  and  $CP_2 \times S^2$ . This will be assumed in the following but should be carefully proven.
3. Locally the spinors of the twistor space  $T(H)$  are tensor products of imbedding spinors and those for  $S^2(M^4) \times S^2(CP_2)$  expressible also as tensor products of spinors for  $S^2(M^4)$  and  $S^2(CP_2)$ . Obviously, the number of spinor components increases by factor  $2 \times 2 = 4$  unless one poses some additional conditions taking care that one has dimensional reduction without the emergence of any new spin like degrees of freedom for which there is no physical evidence. The only possible manner to achieve this is to pose covariant constancy conditions already at the level of twistor spaces  $T(M^4)$  and  $T(CP_2)$  leaving only single spin state in these degrees of freedom.
4. In  $CP_2$  covariant constancy is possible for right-handed neutrino so that  $CP_2$  spinor structure can be taken as a model. In the case of  $CP_2$  spinors covariant constancy is possible for right-handed neutrino and is essentially due to the presence of  $U(1)$  part in spinor connection forced by the fact that the spinor structure does not exist otherwise. Ordinary  $S^2$  spinor connection defined by vielbein exists always. One can however add a coupling to a suitable multiple of Kähler potential satisfying the quantization of magnetic charge (the magnetic flux defined by  $U(1)$  connection is multiple of  $2\pi$  so that its imaginary exponential is unity).

$S^2$  spinor connections must have besides ordinary vielbein part determined by  $S^2$  metric also  $U(1)$  part defined by Kähler form coupled with correct coupling so that the curvature form annihilates the second spin state for both  $S^2(M^4)$  and  $S^2(CP_2)$ .  $U(1)$  part of the spinor curvature is proportional to Kähler form  $J \propto \sin(\theta)d\theta d\phi$  so that this is possible. The vielbein and  $U(1)$  parts of the spinor curvature are proportional Pauli spin matrix  $\sigma_z = (1, 0; 0, -1)/2$  and unit matrix  $(1, 0; 0, 1)$  respectively so that the covariant constancy is possible to satisfy and fixes the spin state uniquely.

5. The covariant derivative for the induced spinors is defined by the sum of projections of spinor gauge potentials for  $T(M^4)$  and  $T(CP_2)$ . With above assumptions the contributions gauge potentials from  $T(M^4)$  and  $T(CP_2)$  separately annihilate single spinor component. As a consequence there are no constraints on the winding numbers  $w_i$ ,  $i = 1, 2$  of the maps  $S^2(X^4) \rightarrow S^2(M^4)$  and  $S^2(X^4) \rightarrow S^2(CP_2)$ . Winding number  $w_i$  corresponds to the imbedding map  $(\Theta_i = \theta, \Phi_i = w_i\phi)$ .
6. If the square of the Kähler form in fiber degrees of freedom gives metric to that its square is metric, one obtains just the area of  $S^2$  from the fiber part of action. This is given by the area  $A = 4\pi\sqrt{2(w_1^2 R_1^2 + w_2^2 R_2^2)}$  since the induced metric is given by  $ds^2 = (R_1^2 + R_2^2)d\theta^2 + (w_1^2 R_1^2 + w_2^2 R_2^2)d\phi^2$  for  $(\Theta_1 = \theta, \Phi = n_1\phi, \Phi_2 = n_2\phi)$ .

## 2.8 Twistor googly problem transforms from a curse to blessing in TGD framework

There was a nice story with title "Michael Atiyah's Imaginative State of Mind" about mathematician Michael Atiyah in Quanta Magazine (see <http://tinyurl.com/jta2va8>). The works of Atiyah have affected profoundly the development of theoretical physics. What was pleasant to hear that Atiyah belongs to those scientists who do not care what others think. As he tells, he can afford this since he has got all possible prizes. This is consoling and encouraging even for those who have not cared what others think and for this reason have not earned any prizes. Nor even a single coin from what they have been busily doing their whole lifetime!

In the beginning of the story "twistor googly problem" was mentioned. I had to refresh my understanding about googly problem. In twistorial description the modes of massless fields (rather than entire massless fields) in space-time are lifted to the modes in its 6-D twistor-space and dynamics reduces to holomorphy. The analog of this takes place also in string models by conformal invariance and in TGD by its extension.

One however encounters what is known as googly problem: one can have twistorial description for circular polarizations with well-defined helicity  $+1/-1$  but not for general polarization states - say linear polarizations, which are superposition of circular polarizations. This reflects itself in the construction of twistorial amplitudes in twistor Grassmann program for gauge fields but rather implicitly: the amplitudes are constructed only for fixed helicity states of scattered particles. For gravitons the situation gets really bad because of non-linearity.

Mathematically the most elegant solution would be to have only  $+1$  or  $-1$  helicity but not their superpositions implying very strong parity breaking and chirality selection. Parity parity breaking occurs in physics but is very small and linear polarizations are certainly possible! The discussion of Penrose with Atiyah has inspired a possible solution to the problem known as "palatial twistor theory" (see <http://tinyurl.com/hr7mh2>). Unfortunately, the article is behind paywall too high for me so that I cannot say anything about it.

What happens to the googly problem in TGD framework? There is twistorialization at space-time level and imbedding space level.

1. One replaces space-time with 4-surface in  $H = M^4 \times CP_2$  and lifts this 4-surface to its 6-D twistor space represented as a 6-surface in 12-D twistor space  $T(H) = T(M^4) \times T(CP_2)$ . The twistor space has Kähler structure only for  $M^4$  and  $CP_2$  so that TGD is unique. This Kähler structure is needed to lift the dynamics of Kähler action to twistor context and the lift leads to the a dramatic increase in the understanding of TGD: in particular, Planck length and cosmological constant with correct sign emerge automatically as dimensional constants besides  $CP_2$  size.
2. Twistorialization at imbedding space level means that spinor modes in  $H$  representing ground states of super-symplectic representations are lifted to spinor modes in  $T(H)$ .  $M^4$  chirality is in TGD framework replaced with H-chirality, and the two chiralities correspond to quarks and leptons. But one cannot superpose quarks and leptons! "Googly problem" is just what the superselection rule preventing superposition of quarks and leptons requires in TGD!

One can look this in more detail.

1. Chiral invariance makes possible for the modes of massless fields to have definite chirality: these modes correspond to holomorphic or antiholomorphic amplitudes in twistor space and holomorphy (antiholomorphy is holomorphy with respect to conjugates of complex coordinates) does not allow their superposition so that massless bosons should have well-defined helicities in conflict with experimental facts. Second basic problem of conformally invariant field theories and of twistor approach relates to the fact that physical particles are massive in 4-D sense. Masslessness in 4-D sense also implies infrared divergences for the scattering amplitudes. Physically natural cutoff is required but would break conformal symmetry.

2. The solution of problems is masslessness in 8-D sense allowing particles to be massive in 4-D sense. Fermions have a well-defined 8-D chirality - they are either quarks or leptons depending on the sign of chirality. 8-D spinors are constructible as superpositions of tensor products of  $M^4$  spinors and of  $CP_2$  spinors with both having well-defined chirality so that tensor product has chiralities  $(\epsilon_1, \epsilon_2)$ ,  $\epsilon_i = \pm 1$ ,  $i = 1, 2$ . H-chirality equals to  $\epsilon = \epsilon_1 \epsilon_2$ . For quarks one has  $\epsilon = 1$  (a convention) and for leptons  $\epsilon = -1$ . For quark states massless in  $M^4$  sense one has either  $(\epsilon_1, \epsilon_2) = (1, 1)$  or  $(\epsilon_1, \epsilon_2) = (-1, -1)$  and for massive states superposition of these. For leptons one has either  $(\epsilon_1, \epsilon_2) = (1, -1)$  or  $(\epsilon_1, \epsilon_2) = (-1, 1)$  in massless case and superposition of these in massive case.
3. The twistorial lift to  $T(M^4) \times T(CP_2)$  of the ground states of super-symplectic representations represented in terms of tensor products formed from H-spinor modes involves only quark and lepton type spinor modes with well-defined H-chirality. Superpositions of amplitudes in which different  $M^4$  helicities appear but  $M^4$  chirality is always paired with completely correlating  $CP_2$  chirality to give either  $\epsilon = 1$  or  $\epsilon = -1$ . One has never a superposition of of different chiralities in either  $M^4$  or  $CP_2$  tensor factor. I see no reason forbidding this kind of mixing of holomorphicities and this is enough to avoid googly problem. Linear polarizations and massive states represent states with entanglement between  $M^4$  and  $CP_2$  degrees of freedom. For massless and circularly polarized states the entanglement is absent.
4. This has interesting implications for the massivation. Higgs field cannot be scalar in 8-D sense since this would make particles massive in 8-D sense and separate conservation of  $B$  and  $L$  would be lost. Theory would also contain a dimensional coupling. TGD counterpart of Higgs boson is actually  $CP_2$  vector, and one can say that gauge bosons and Higgs combine to form 8-D vector. This correctly predicts the quantum numbers of Higgs. Ordinary massivation by constant vacuum expectation value of vector Higgs is not an attractive idea since no covariantly constant  $CP_2$  vector field exists so that Higgsy massivation is not promising except at QFT limit of TGD formulated in  $M^4$ . p-Adic thermodynamics gives rise to 4-D massivation but keeps particles massless in 8-D sense. It also leads to powerful and correct predictions in terms of p-adic length scale hypothesis.

Anonymous reader gave me a link to the paper of Penrose about palatial twistor theory and this inspired further more detailed considerations of googly problem.

1. After the first reading I must say that I could not understand how the proposed elimination of conjugate twistor by quantization of twistors solves the googly problem, which means that both helicities are present (twistor  $Z$  and its conjugate) in linearly polarized classical modes so that holomorphy is broken classically.
2. I am also very skeptic about quantizing of either space-time coordinates or twistor space coordinates. To me quantization is natural only for linear objects like spinors. For bosonic objects one must go to higher abstraction level and replace superpositions in space-time with superpositions in field space. Construction of "World of Classical Worlds" (WCW) in TGD means just this.
3. One could however think that circular polarizations are fundamental and quantal linear combination of the states carrying circularly polarized modes give rise to linear and elliptic polarizations. Linear combination would be possible only at the level of field space (WCW in TGD), not for classical fields in space-time. If so, then the elimination of conjugate of  $Z$  by quantization suggested by Penrose would work.
4. Unfortunately, Maxwell's equations allow classically linear polarisations! In order to achieve classical-quantum consistency, one should modify classical Maxwell's equations somehow so that linear polarizations are not possible. Googly problem is still there!

What about TGD?

1. Massless extremals representing massless modes are very "quantal": they cannot be superposed classically unless both momentum and polarisation directions for them (they can depend space-time point) are exactly parallel. Optimist would guess that the classical local classical polarisations are circular. No, they are linear! Superposition of classical linear polarizations at the level of WCW can give rise to local linear but not local circular polarization! Something more is needed.
2. The only sensible conclusion is that only gauge boson quanta (not classical modes) represented as pairs of fundamental fermion and antifermion in TGD framework can have circular polarization! And indeed, massless bosons - in fact, all elementary particles- are constructed from fundamental fermions and they allow only two  $M^4$ ,  $CP_2$  and  $M^4 \times CP_2$  helicities/-chiralities analogous to circular polarisations. B and L conservation would transform googly problem to a superselection rule as already described.

To sum up, both the extreme non-linearity of Kähler action, the representability of all elementary particles in terms of fundamental fermions and antifermions, and the generalization of conserved  $M^4$  chirality to conservation of H-chirality would be essential for solving the googly problem in TGD framework.

### 3 Surprise: twistorial dynamics does not reduce to a trivial reformulation of the dynamics of Kähler action

I have thought that twistorialization classically means only an alternative formulation of TGD. This is definitely not the case as the explicit study demonstrated. Twistor formulation of TGD is in terms of 6-D twistor spaces  $T(X^4)$  of space-time surfaces  $X^4 \subset M^4 \times CP_2$  in 12-dimensional product  $T = T(M^4) \times T(CP_2)$  of 6-D twistor spaces of  $T(M^4)$  of  $M^4$  and  $T(CP_2)$  of  $CP_2$ . The induced Kähler form in  $X^4$  defines the quaternionic imaginary unit defining twistor structure: how stupid that I realized it only now! I experienced during single night many other "How stupid I have been" experiences.

Classical dynamics is determined by 6-D variant of Kähler action with coefficient  $1/L^2$  having dimensions of inverse length squared. Since twistor space is bundle, a dimensional reduction of 6-D Kähler action to 4-D Kähler action plus a term analogous to cosmological term - space-time volume - takes place so that dynamics reduces to 4-D dynamics also now. Here one must be careful: this happens provided the radius of  $S^2$  associated with  $X^4$  does not depend on point of  $X^4$ . The emergence of cosmological term was however completely unexpected: again "How stupid I have been" experience. The scales of the spheres and the condition that the 6-D action is dimensionless bring in 3 fundamental length scales!

#### 3.1 New scales emerge

The twistorial dynamics gives to several new scales with rather obvious interpretation. The new fundamental constants that emerge are the radii of the spheres associated with  $T(M^4)$  and  $T(CP_2)$ . The radius of the sphere associated with  $X^4$  is not a fundamental constant but determined by the induced metric. By above argument the fiber is sphere for both Euclidian signature and Minkowskian signatures.

1. For  $CP_2$  twistor space the radius of  $S^2(CP_2)$  must be apart from numerical constant equal to  $CP_2$  radius  $R$ . For  $S^2(M^4)$  one can consider two options. The first option is that also now the radius for  $S^2(M^4)$  equals to  $R(M^4) = R$  so that Planck length would not emerge from fundamental theory classically as assumed hitherto. Second imaginable option is that it does and one has  $R(M^4) = l_P$ .
2. If the signature of  $S^2(M^4)$  is  $(-1, -1)$  both Minkowskian and Euclidian regions have  $S^2(X^4)$  with the same signature  $(-1, -1)$ . The radius  $R_D$  of  $S^2(X^4)$  is dynamically determined.

Recall first how the cosmological constant emerges from TGD framework. The key point is that the 6-D Kähler action contains two terms.

1. The first term is essentially the ordinary Kähler action multiplied by the area of  $S^2(X^4)$  which is compensated by the length scale, which can be taken to be the area  $4\pi R^2(M^4)$  of  $S^2(M^4)$ . This makes sense for winding numbers  $(w_1, w_2) = (1, 0)$  meaning that  $S^2(CP_2)$  is effectively absent but  $S^2(M^4)$  is present.
2. Second term is the analog of Kähler action assignable assignable to the projection of  $S^2(M^4)$  Kähler form. The corresponding Kähler coupling strength  $\alpha_K(M^4)$  is huge - so huge that one has  $\alpha_K(M^4)4\pi R^2(M^4) \equiv L^2$ , where  $1/L^2$  is of the order of cosmological constant and thus of the order of the size of the recent Universe.  $\alpha_K(M^4)$  is also analogous to critical temperature and the earlier hypothesis that the values of  $L$  correspond to p-adic length scales implies that the values of come as  $\alpha_K(M^4) \propto p \simeq 2^k$ ,  $p$  prime,  $k$  prime.
3. One can get an estimate for the relative magnitude of the Kähler action  $S(CP_2) = \pi/8\alpha_K$  assignable to  $CP_2$  type vacuum extremal and the corresponding cosmological term. The magnitude of the volume term is of order  $1/4\pi\alpha_K(M^4)$  with  $\alpha_K(M^4)$  given by  $\alpha_K(M^4) = L^2/4\pi R^2(M^4)$ . The sequel the magnitude of  $L$  is estimated to be  $L = (2^{3/2}\pi l_P/R_D) \times R_U$ , where  $R_U$  is the recent size of the Universe. This estimate follows from the identification of the volume term as cosmological constant term.

For  $R_D = R_M = l_P$  this gives  $\alpha_K(M^4) = 2\pi(R_U/l_P)^2 \sim 2 \times 10^{18}$ . For  $\alpha_K \simeq 1/137$  the ratio of the two terms is of order  $10^{-20}$ . The cosmological terms is completely negligible in elementary particle scales. For vacuum extremals the situation changes and the overall effect is presumably the transformation of 4-D spin glass degeneracy so that the potentials wells in the analog spin glass energy landscape do not correspond to vacuum extremal anymore and perturbation theory around them is in principle possible. The huge value of  $\alpha_K(M^4)$  implies that the system corresponds mathematically to an extremely strongly interacting system so that perturbation theory fails to converge. The geometry of "World of Classical Worlds" (WCW) provides the needed non-perturbative approach and leads to strong form of holography.

4. One could argue that the Kähler form assignable to  $M^4$  cannot contribute to the action since it does not contribute to spinor connection of  $M^4$  - an assumption that can be challenged. For canonically imbedded  $M^4$  self-duality implies that this contribution to action vanishes. For vacuum extremals of ordinary Kähler action the contribution to the action density is proportional to the  $CP_2$  part of induced metric and to  $1/\alpha_K(M^4)$ , and therefore extremely small.

The breaking of Lorentz invariance can be seen as a possible problem for the induced spinor fields coupling to the self-dual Kähler potential. This corresponds to coupling to constant magnetic field and constant electric field, which are duals of each other. This would give rise to the analogs of cyclotron energy states in transversal directions and to the analogs of states in constant electric field in longitudinal directions. Could this extremely small effect serve as a seed for the generation of Kähler magnetic flux tubes carrying longitudinal electric fields in various scales? Note also that the value of  $\alpha_K(M^4)$  is predicted to decrease as p-adic length scale so that the effect would be larger in early cosmology and in short length scales.

Hence one can consider the possibility that the action is just the sum of full 6-D Kähler actions assignable to  $T(M^4)$  and  $T(CP_2)$  but with different values of  $\alpha_K$  if one has  $(w_1, w_2) = (n, 0)$ . Also other  $w_2 \neq 0$  is possible but corresponds to gigantic cosmological constant.

Given the parameter  $L^2$  as it is defined above, one can deduce an expression for cosmological constant  $\Lambda$  and show that it is positive.

1. 6-D Kähler action has dimensions of length squared and one must scale it by a dimensional constant: call it  $1/L^2$ .  $L$  is a fundamental scale and in dimensional reduction it gives rise to cosmological constant. Cosmological constant  $\Lambda$  is defined in terms of vacuum energy density as  $\Lambda = 8\pi G\rho_{vac}$  can have two interpretations.  $\Lambda$  can correspond to a modification of Einstein-Hilbert action or - as

now - to an additional term in the action for matter. In the latter case positive  $\Lambda$  means negative pressure explaining the observed accelerating expansion. It is actually easy to deduce the sign of  $\Lambda$ .  $1/L^2$  multiplies both Kähler action -  $F^{ij}F_{ij}$  ( $\propto E^2 - B^2$  in Minkowskian signature). The energy density is positive. For Kähler action the sign of the multiplier must be positive so that  $1/L^2$  is positive. The volume term is fiber space part of action having same form as Kähler action. It gives a positive contribution to the energy density and negative contribution to the pressure.

In  $\Lambda = 8\pi G\rho_{vac}$  one would have  $\rho_{vac} = \pi/L^2 R_D^2$  as integral of the  $-F^{ij}F_{ij}$  over  $S^2$  given the  $\pi/R_D^2$  (no guarantee about correctness of numerical constants). This gives  $\Lambda = 8\pi^2 G/L^2 R_D^2$ .  $\Lambda$  is positive and the sign is same as as required by accelerated cosmic expansion. Note that super string models predict wrong sign for  $\Lambda$ .  $\Lambda$  is also dynamical since it depends on  $R_D$ , which is dynamical. One has  $1/L^2 = k\Lambda$ ,  $k = 8\pi^2 G/R_D^2$  apart from numerical factors.

The value of  $L$  of deduced from Euclidian and Minkowskian regions in this formal manner need not be same. Since the GRT limit of TGD describes space-time sheets with Minkowskian signature, the formula seems to be applicable only in Minkowskian regions. Again one can argue that one cannot exclude Euclidian space-time sheets of even macroscopic size and blackholes and even ordinary concept matter would represent this kind of structures.

2.  $L$  is not a size scale of any fundamental geometric object. This suggests that  $L$  is analogous to  $\alpha_K$  and has value spectrum dictated by p-adic length scale hypothesis. In fact, one can introduce the ratio of  $\epsilon = R^2/L^2$  as a dimensionless parameter analogous to coupling strength what it indeed is in field equations. If so,  $L$  could have different values in Minkowskian and Euclidian regions.
3. I have earlier proposed that  $R_U \equiv 1/\sqrt{1/\Lambda}$  is essentially the p-adic length scale  $L_p \propto \sqrt{p} = 2^{k/2}$ ,  $p \simeq 2^k$ ,  $k$  prime, characterizing the cosmology at given time and satisfies  $R_U \propto a$  meaning that vacuum energy density is piecewise constant but on the average decreases as  $1/a^2$ ,  $a$  cosmic time defined by light-cone proper time. A more natural hypothesis is that  $L$  satisfies this condition and in turn implies similar behavior or  $R_U$ . p-Adic length scales would be the critical values of  $L$  so that also p-adic length scale hypothesis would emerge from quantum critical dynamics! This conforms with the hypothesis about the value spectrum of  $\alpha_K$  labelled in the same manner [11].
4. At GRT limit the magnetic energy of the flux tubes gives rise to an average contribution to energy momentum tensor, which effectively corresponds to negative pressure for which the expansion of the Universe accelerates. It would seem that both contributions could explain accelerating expansion. If the dynamics for Kähler action and volume term are coupled, one would expect same orders of magnitude for negative pressure and energy density - kind of equipartition of energy.

Consider first the basic scales emerging also from GRT picture.  $R_U \sim \sqrt{1/\Lambda} \sim 10^{26}$  m = 10 Gly is not far from the recent size of the Universe defined as  $c \times t \sim 13.8$  Gly. The derived size scale  $L_1 \equiv (R_U \times l_P)^{1/2}$  is of the order of  $L_1 = .5 \times 10^{-4}$  meters, the size of neuron. Perhaps this is not an accident. To make life of the reader easier I have collected the basic numbers to the following table.

$$\begin{aligned}
 m(CP_2) &\simeq 5.7 \times 10^{14} \text{ GeV} , & m_P &= 2.435 \times 10^{18} \text{ GeV} , & \frac{R(CP_2)}{l_P} &\simeq 4.1 \times 10^3 , \\
 R_U &= 10 \text{ Gy} , & t &= 13.8 \text{ Gy} , & L_1 &= \sqrt{l_P R_U} = .5 \times 10^{-4} \text{ m} .
 \end{aligned}
 \tag{3.1}$$

Let us consider now some quantitative estimates.  $R(X^4)$  depends on homotopy equivalence classes of the maps from  $S^2(X^4) \rightarrow S^2(M^4)$  and  $S^2(X^4) \rightarrow S^2(CP_2)$  - that is winding numbers  $w_i$ ,  $i = 1, 2$  for these maps. The simplest situations correspond to the winding numbers  $(w_1, w_2) = (1, 0)$  and  $(w_1, w_2) = (0, 1)$ . For  $(w_1, w_2) = (1, 0)$   $M^4$  contribution to the metric of  $S^2(X^4)$  dominates and one has  $R(X^4) \simeq R(M^4)$ .

For  $R(M^4) = l_P$  so Planck length would define a fundamental length and Planck mass and Newton's constant would be quantal parameters. For  $(w_1, w_2) = (0, 1)$  the radius of sphere would satisfy  $R_D \simeq R$  ( $CP_2$  size): now also Planck length would be quantal parameter.

Consider next additional scales emerging from TGD picture.

1. One has  $L = (2^{3/2}\pi l_P/R_D) \times R_U$ . In Minkowskian regions with  $R_D = l_P$  this would give  $L = 8.9 \times R_U$ : there is no obvious interpretation for this number in recent cosmology. For ( $R_D = R$ ) one obtains the estimate  $L = 29$  Mly. The size scale of large voids varies from about 36 Mly to 450 Mly (see [https://en.wikipedia.org/wiki/Void\\_\(astronomy\)](https://en.wikipedia.org/wiki/Void_(astronomy))).
2. Consider next the derived size scale  $L_2 = (L \times l_P)^{1/2} = \sqrt{L/R_U} \times L_1 = \sqrt{2^{3/2}\pi l_P/R_D} \times L_1$ . For  $R_D = l_P$  one has  $L_2 \simeq 3L_1$ . For  $R_D = R$  making sense in Euclidian regions, this is of the order of size of neutrino Compton length:  $3 \mu\text{m}$ , the size of cellular nucleus and rather near to the p-adic length scale  $L(167) = 2.6$  m, corresponds to the largest miracle Gaussian Mersennes associated with  $k = 151, 157, 163, 167$  defining length scales in the range between cell membrane thickness and the size of cellular nucleus. Perhaps these are co-incidences are not accidental. Biology is something so fundamental that fundamental length scale of biology should appear in the fundamental physics.

The formulas and predictions for different options are summarized by the following table.

$$\begin{array}{lll}
 \text{Option} & L = \frac{2^{3/2}\pi l_P}{R_D} \times R_U & L_2 = \sqrt{L l_P} = \sqrt{\frac{2^{3/2}\pi l_P}{R_D}} \times L_1 \\
 R_D = R & , \quad 29 \text{ Mly} & , \quad \simeq 3 \mu\text{m} \\
 R_D = l_P & , \quad 8.9 R_U & , \quad \simeq 3L_1 = 1.5 \times 10^{-4} \text{ m}
 \end{array} \tag{3.2}$$

In the case of  $M^4$  the radius of  $S^2$  cannot be fixed it remains unclear whether Planck length scale is fundamental constant or whether it emerges.

### 3.2 Estimate for the cosmic evolution of $R_D$

One can actually get estimate for the evolution of  $R_D$  as function of cosmic time if one accepts Friedman cosmology as an approximation of TGD cosmology.

1. Assume critical mass density so that one has

$$\rho_{cr} = \frac{3H^2}{8\pi G} .$$

2. Assume that the contribution of cosmological constant term to the mass mass density dominates. This gives  $\rho \simeq \rho_{vac} = \Lambda/8\pi G$ . From  $\rho_{cr} = \rho_{vac}$  one obtains

$$\Lambda = 3H^2 .$$

3. From Friedman equations one has  $H^2 = ((da/dt)/a)^2$ , where  $a$  corresponds to light-cone proper time and  $t$  to cosmic time defined as proper time along geodesic lines of space-time surface approximated as Friedmann cosmology. One has

$$\Lambda = \frac{3}{g_{aa}a^2}$$

in Robertson-Walker cosmology with  $ds^2 = g_{aa}da^2 - a^2d\sigma_3^2$ .

4. Combining this equations with the TGD based equation

$$\Lambda = \frac{8\pi^2 G}{L^2 R_D^2}$$

one obtains

$$\frac{8\pi^2 G}{L^2 R_D^2} = \frac{3}{g_{aa} a^2} . \quad (3.3)$$

5. Assume that quantum criticality applies so that  $L$  has spectrum given by p-adic length scale hypothesis so that one discrete p-adic length scale evolution for the values of  $L$ . There are two options to consider depending on whether p-adic length scales are assigned with light-cone proper time  $a$  or with cosmic time  $t$

$$T = a \text{ (Option I) } , \quad T = t \text{ (Option II)} \quad (3.4)$$

Both options give the same general formula for the p-adic evolution of  $L(k)$  but with different interpretation of  $T(k)$ .

$$\frac{L(k)}{L_{now}} = \frac{T(k)}{T_{now}} , \quad T(k) = L(k) = 2^{(k-151)/2} \times L(151) , \quad L(151) \simeq 10 \text{ nm} . \quad (3.5)$$

Here  $T(k)$  is assumed to correspond to primary p-adic length scale. An alternative - less plausible - option is that  $T(k)$  corresponds to secondary p-adic length scale  $L_2(k) = 2^{k/2} L(k)$  so that  $T(k)$  would correspond to the size scale of causal diamond. In any case one has  $L \propto L(k)$ . One has a discretized version of smooth evolution

$$L(a) = L_{now} \times \frac{T}{T_{now}} . \quad (3.6)$$

6. Feeding into this to Eq. 3.3 one obtains an expression for  $R_D(a)$

$$\frac{R_D}{l_P} = \left(\frac{8}{3}\right)^{1/2} \pi \times \frac{a}{L(a)} \times g_{aa}^{1/2} . \quad (3.7)$$

Unless the dependences on cosmic time compensate each other,  $R_D$  is dynamical and becomes very small at very early times since  $g_{aa}$  becomes very small.  $R(M^4) = l_P$  however poses a lower boundary since either of the maps  $S^2(X^4) \rightarrow S^2(M^4)$  and  $S^2(X^4) \rightarrow S^2(CP_2)$  must be homotopically non-trivial. For  $R(M^4) = l_P$  one would obtain  $R_D/l_P = 1$  at this limit giving also lower bound for  $g_{aa}$ . For  $T = t$  option  $a/L(a)$  becomes large and  $g_{aa}$  small.

As a matter of fact, in very early cosmic string dominated cosmology  $g_{aa}$  would be extremely small constant [5]. In late cosmology  $g_{aa} \rightarrow 1$  holds true and one obtains at this limit

$$\frac{R_D(now)}{l_P} = \left(\frac{8}{3}\right)^{1/2} \pi \times \frac{a_{now}}{L_{now}} \times l_P \simeq 4.4 \frac{a_{now}}{L_{now}} . \quad (3.8)$$

7. For  $T = t$  option  $R_D/l_P$  remains constant during both matter dominated cosmology, radiation dominated cosmology, and string dominated cosmology since one has  $a \propto t^n$  with  $n = 1/2$  during radiation dominated era,  $n = 2/3$  during matter dominated era, and  $n = 1$  during string dominated era [5]. This gives

$$\frac{R_D}{l_P} = \left(\frac{8}{3}\right)^{1/2} \pi \times \frac{a}{t} \sqrt{g_{aa}} \frac{t(\text{end})}{L(\text{end})} = \left(\frac{8}{3}\right)^{1/2} \frac{\pi}{n} \frac{t(\text{end})}{L(\text{end})} .$$

Here "end" refers the end of the string or radiation dominated period or to the recent time in the case of matter dominated era. The value of  $n$  would have evolved as  $R_D/l_P \propto (1/n)(t_{\text{end}}/L_{\text{end}})$ ,  $n \in \{1, 3/2, 2\}$ . During radiation dominated cosmology  $R_D \propto a^{1/2}$  holds true. The value of  $R_D$  would be very nearly equal to  $R(M^4)$  and  $R(M^4)$  would be of the same order of magnitude as Planck length. In matter dominated cosmology would have  $R_D \simeq 2.2(t(\text{now})/L(\text{now})) \times l_P$ .

8. For  $R_D(\text{now}) = l_P$  one would have

$$\frac{L_{\text{now}}}{a_{\text{now}}} = \left(\frac{8}{3}\right)^{1/2} \pi \simeq 4.4 .$$

In matter dominated cosmology  $g_{aa} = 1$  gives  $t_{\text{now}} = (2/3) \times a_{\text{now}}$  so that predictions differ only by this factor for options I and II. The winding number for the map  $S^2(X^4) \rightarrow S^2(CP_2)$  must clearly vanish since otherwise the radius would be of order  $R$ .

9. For  $R_D(\text{now}) = R$  one would obtain

$$\frac{a_{\text{now}}}{L_{\text{now}}} = \left(\frac{8}{3}\right)^{1/2} \times \frac{R}{l_P} \simeq 2.1 \times 10^4 .$$

One has  $L_{\text{now}} = 10^6$  ly: this is roughly the average distance scale between galaxies. The size of Milky Way is in the range  $1 - 1.8 \times 10^5$  ly and of an order of magnitude smaller.

10. An interesting possibility is that  $R_D(a)$  evolves from  $R_D \sim R(M^4) \sim l_P$  to  $R_D \sim R$ . This could happen if the winding number pair  $(w_1, w_2) = (1, 0)$  transforms to  $(w_1, w_2) = (0, 1)$  during transition to from radiation (string) dominance to matter (radiation) dominance.  $R_D/l_P$  radiation dominated cosmology would be related by a factor

$$\frac{R_D(\text{rad})}{R_D(\text{mat})} = (3/4) \frac{t(\text{rad}, \text{end})}{L(\text{rad}, \text{end})} \times \frac{L(\text{now})}{t(\text{now})}$$

to that in matter dominated cosmology. Similar factor would relate the values of  $R_D/l_P$  in string dominated and radiation dominated cosmologies. The condition  $R_D(\text{rad})/R_D(\text{mat}) = l_P/R$  expressing the transformation of winding numbers would give

$$\frac{L(\text{now})}{L(\text{rad}, \text{end})} = \frac{4 l_P}{3 R} \frac{t(\text{now})}{t(\text{rad}, \text{end})} .$$

One has  $t(\text{now})/t(\text{rad}, \text{end}) \simeq .5 \times 10^6$  and  $l_P/R = 2.5 \times 10^{-4}$  giving  $L(\text{now})/L(\text{rad}, \text{end}) \simeq 125$ , which happens to be near fine structure constant.

11. For the twistorial lifts of space-time surfaces for which cosmological constant has a reasonable value, the winding numbers are equal to  $(w_1, w_2) = (n, 0)$  so that  $R_D = \sqrt{n}R(S^2(M^4))$  holds true in good approximation. This conforms with the observed constancy of  $R_D$  during various cosmological eras, and would suggest that the ratio  $\frac{t(\text{end})}{L(\text{end})}$  characterizing these periods is same for all periods. This determines the evolution for the values of  $\alpha_K(M^4)$ .

$R(M^4) \sim l_P$  seems rather plausible option so that Planck length would be fundamental classical length scale emerging naturally in twistor approach. Cosmological constant would be coupling constant like parameter with a spectrum of critical values given by p-adic length scales.

### 3.3 What about extremals of the dimensionally reduced 6-D Kähler action?

It seems that the basic wisdom about extremals of Kähler action remains unaffected and the motivations for WCW are not lost. What is new is that the removal of vacuum degeneracy is forced by twistorial action.

1. All extremals, which are minimal surfaces remain extremals. In fact, all the known extremals except vacuum extremals. For minimal surfaces the dynamics of the volume term and 4-D Kähler action separate and field equations for them are separately satisfied. The vacuum degeneracy motivating the introduction of WCW is preserved. The induced Kähler form vanishes for vacuum extremals and the imaginary unit of twistor space is ill-defined. Hence vacuum extremals cannot belong to WCW. This correspond to the vanishing of WCW metric for vacuum extremals.
2. For non-minimal surfaces Kähler coupling strength does not disappear from the field equations and appears as a genuine coupling very much like in classical field theories. Minimal surface equations are a generalization of wave equation and Kähler action would define analogs of source terms. Field equations would state that the total isometry currents are conserved. It is not clear whether other than minimal surfaces are possible, I have even conjectured that all preferred extremals are always minimal surfaces having the property that being holomorphic they are almost universal extremals for general coordinate invariant actions.
3. Thermodynamical analogy might help in the attempts to interpret. Quantum TGD in zero energy ontology (ZEO) corresponds formally to a complex square root of thermodynamics. Kähler action can be identified as a complexified analog of free energy. Complexification follows both from the fact that  $\sqrt{g}$  is real/imaginary in Euclidian/Minkowskian space-time regions. Complex values are also implied by the proposed identification of the values of Kähler coupling strength in terms of zeros and pole of Riemann zeta in turn identifiable as poles of the so called fermionic zeta defining number theoretic partition function for fermions [8, 11, 12]. The thermodynamical for Kähler action with volume term is Gibbs free energy  $G = F - TS = E - TS + PV$  playing key role in chemistry.
4. The boundary conditions at the ends of space-time surfaces at boundaries of CD generalize appropriately and symmetries of WCW remain as such. At light-like boundaries between Minkowskian and Euclidian regions boundary conditions must be generalized. In Minkowskian regions volume can be very large but only the Euclidian regions contribute to Kähler function so that vacuum functional can be non-vanishing for arbitrarily large space-time surfaces since exponent of Minkowskian Kähler action is a phase factor.
5. One can worry about almost topological QFT property. Although Kähler action from Minkowskian regions at least would reduce to Chern-Simons terms with rather general assumptions about preferred extremals, the extremely small cosmological term does not. Could one say that cosmological constant term is responsible for "almost"?

It is interesting that the volume of manifold serves in algebraic geometry as topological invariant for hyperbolic manifolds, which look locally like hyperbolic spaces  $H_n = SO(n, 1)/SO(n)$  [1, 7]. See also the article "Volumes of hyperbolic manifolds and mixed Tate motives" (see <http://arxiv.org/abs/alg-geom/9601021>). Now one would have  $n = 4$ . It is probably too much to hope that space-time surfaces would be hyperbolic manifolds. In any case, by the extreme uniqueness of the preferred extremal property expressed by strong form of holography the volume of space-time surface could also now serve as topological invariant in some sense as I have earlier proposed. What is intriguing is that  $AdS_n$  appearing in AdS/CFT correspondence is Lorentzian analogue  $H_n$ .

6.  $\alpha(M^4)$  is extremely large so that there is no hope of quantum perturbation theory around canonically imbedded  $M^4$  although the propagator for  $CP_2$  coordinate exists. In the new framework WCW can be seen as a solution to how to construct non-perturbative quantum TGD.

To sum up, I have the feeling that the final formulation of TGD has now emerged and it is clear that TGD is indeed a quantum theory of gravitation allowing to understand standard model symmetries. The existence of twistorial formulation is all that is needed to fix the theory completely. It makes possible gravitation and predicts standard model symmetries. This cannot be said about any competitor of TGD.

## 4 Basic principles behind construction of amplitudes

Basic principles of the construction summarized in this section could be seen as axioms trying to abstract the essentials. The explicit construction of amplitudes is too heavy challenge at this stage and at least for me.

### 4.1 Imbedding space is twistorially unique

It took roughly 36 years to learn that  $M^4$  and  $CP_2$  are twistorially unique.

1. As already explained,  $M^4$  and  $CP_2$  are unique 4-manifolds in the sense that both allow twistor space with Kähler structure: Kähler structure is the crucial concept as one might guess from the fact that the projection of Kähler form naturally defines the preferred quaternionic imaginary unit defining the twistor structure for space-time surface. Both  $M^4$  and its Euclidian variant  $E^4$  allow twistor space and the twistor space of  $M^4$  is Minkowskian variant  $T(M^4) = SU(2, 2)/SU(2, 1) \times U(1)$  of 6-D twistor space  $CP_3 = SU(4)/SU(3) \times U(1)$  of  $E^4$ . The twistor space of  $CP_2$  is 6-D  $T(CP_2) = SU(3)/U(1) \times U(1)$ , the space for the choices of quantization axes of color hypercharge and isospin.
2. This leads to a proposal for the formulation of TGD in which space-time surfaces  $X^4$  in  $H$  are lifted to twistor spaces  $X^6$ , which are sphere bundles over  $X^4$  and such that they are surfaces in 12-D product space  $T(M^4) \times T(CP_2)$  such the twistor structure of  $X^4$  are in some sense induced from that of  $T(M^4) \times T(CP_2)$ . What is nice in this formulation is that one can use all the machinery of algebraic geometry so powerful in superstring theory (Calabi-Yau manifolds). It has been already described how this approach leads to a profound understanding of the relationship between TGD and GRT. Planck length emerges whereas fundamental constant as also cosmological constant emerges dynamically from the length scale parameter appearing in 6-D Kähler action. One can say, that twistor extension is absolutely essential for really understanding the gravitational interactions although the modification of Kähler action is extremely small due to the huge value of length scale defined by cosmological constant.
3. Masslessness (masslessness in complex sense for virtual particles in twistorialization) is essential condition for twistorialization. In TGD massless is masslessness in 8-D sense for the representations of superconformal algebras. This suggests that 8-D variant of twistors makes sense. 8-dimensionality indeed allows octonionic structure in the tangent space of imbedding space. One can also define octonionic gamma matrices and this allows a possible generalization of 4-D twistors to 8-D ones using generalization of sigma matrices representing quaternionic units to octonionic sigma "matrices" essential for the notion of twistors. These octonion units do not of course allow matrix representation unless one restricts to units in some quaternionic subspace of octonions. Space-time surfaces would be associative and thus have quaternionic tangent space at each point satisfying some additional conditions.

## 4.2 Strong form of holography

Strong form of holography (SH) following from general coordinate invariance (GCI) for space-times as surfaces states that the data assignable to string world sheets and partonic 2-surfaces allows to code for scattering amplitudes. The boundaries of string world sheets at the space-like 3-surfaces defining the ends of space-time surfaces at boundaries of causal diamonds (CDs) and the fermionic lines along light-like orbits of partonic 2-surfaces representing lines of generalized Feynman diagrams become the basic elements in the generalization of twistor diagrams (I will not use the attribute "Feynman" in precise sense, one could replace it with "twistor" or even drop away). One can assign fermionic lines massless in 8-D sense to flux tubes, which can also be braided. One obtains a fractal hierarchy of braids with strands, which are braids themselves. At the lowest level one has braids for which fermionic lines are braided. This fractal hierarchy is unavoidable and means generalization of the ordinary Feynman diagram. I have considered some implications of this hierarchy in [13].

## 4.3 The existence of WCW demands maximal symmetries

Quantum TGD reduces to the construction of Kähler geometry of infinite-D WCW, of associated spinor structure, and of modes of WCW spinor fields which are purely classical entities and quantum jump remains the only genuinely quantal element of quantum TGD. Quantization without quantization, would Wheeler say.

By its infinite-dimensionality, the mere mathematical existence of the Kähler geometry of WCW requires maximal isometries. Physics is completely fixed by the mere condition that its mathematical description exists. Super-symplectic and other symmetries of WCW are in decisive role. These symmetry algebras have conformal structure and generalize and extend the conformal symmetries of string models (Kac-Moody algebras in particular). These symmetries give also rise to the hierarchy of Planck constants. The super-symplectic symmetries extend to a Yangian algebra, whose generators are polylocal in the sense that they involve products of generators associated with different partonic surfaces. These symmetries leave scattering amplitudes invariant. This is an immensely powerful constraint, which remains to be understood.

## 4.4 Quantum criticality

Quantum criticality (QC) of TGD Universe is a further principle. QC implies that Kähler coupling strength is mathematically analogous to critical temperature and has a discrete spectrum. Coupling constant evolution is replaced with a discrete evolution as function of p-adic length scale: sequence of jumps from criticality to a more refined criticality or vice versa (in spin glass energy landscape you at bottom of well containing smaller wells and you go to the bottom of smaller well). This implies that either all radiative corrections (loops) sum up to zero (QFT limit) or that diagrams containing loops correspond to the same scattering amplitude as tree diagrams so that loops can be eliminated by transforming them to arbitrary small ones and snipping away moving the end points of internal lines along the lines of diagram (fundamental description).

Quantum criticality at the level of super-conformal symmetries leads to the hierarchy of Planck constants  $h_{eff} = n \times h$  labelling a hierarchy of sub-algebras of super-symplectic and other conformal algebras isomorphic to the full algebra. Physical interpretation is in terms of dark matter hierarchy. One has conformal symmetry breaking without conformal symmetry breaking as Wheeler would put it.

## 4.5 Physics as generalized number theory, number theoretical universality

Physics as generalized number theory vision has important implications. Adelic physics is one of them. Adelic physics implied by number theoretic universality (NTU) requires that physics in real and various

p-adic numbers fields and their extensions can be obtained from the physics in their intersection corresponding to an extension of rationals. This is also enormously powerful condition and the success of p-adic length scale hypothesis and p-adic mass calculations can be understood in the adelic context.

In TGD inspired theory of consciousness various p-adic physics serve as correlates of cognition and p-adic space-time sheets can be seen as cognitive representations, "thought bubbles". NTU is closely related to SH. String world sheets and partonic 2-surfaces with parameters (WCW coordinates) characterizing them in the intersection of rationals can be continued to space-time surfaces by preferred extremal property but not always. In p-adic context the fact that p-adic integration constants depend on finite number of binary digits makes the continuation easy but in real context this need not be possible always. It is always possible to imagine something but not always actualize it!

#### 4.6 Scattering diagrams as computations

Quantum criticality as possibility to eliminate loops has a number theoretic interpretation. Generalized Feynman diagram can be interpreted as a representation of a computation connecting given set  $X$  of algebraic objects to second set  $Y$  of them (initial and final states in scattering) (trivial example:  $X = \{3, 4\} \rightarrow 3 \times 4 = 12 \rightarrow 2 \times 6 \rightarrow \{2, 6\} = Y$ ). The 3-vertices ( $a \times b = c$ ) and their time-reversals represent algebraic product and co-product.

There is a huge symmetry: all diagrams representing computation connecting given  $X$  and  $Y$  must produce the same amplitude and there must exist minimal computation. The task of finding this computation is like finding the simplest representation for the formula  $X=Y$  and the noble purpose of math teachers is that we should learn to find it during our school days. This generalizes the duality symmetry of old fashioned string models: one can transform any diagram to a tree diagram without loops. This corresponds to quantum criticality in TGD: coupling constants do not evolve. The evolution is actually there but discrete and corresponds to infinite number critical values for Kahler coupling strength analogous to temperature.

#### 4.7 Reduction of diagrams with loops to braided tree-diagrams

1. In TGD pointlike particles are replaced with 3-surfaces and by SH by partonic 2-surfaces. The important implication of 3-dimensionality is braiding. The fermionic lines inside light-like orbits of partonic 2-surfaces can be knotted and linked - that is braided (this is dynamical braiding analogous to dance). Also the fermionic strings connecting partonic 2-surfaces at space-like 3-surfaces at boundaries of causal diamonds (CDs) are braided (space-like braiding).

Therefore ordinary Feynman diagrams are not enough and one must allow braiding for tree diagrams. One can also imagine of starting from braids and allowing 3-vertices for their strands (product and co-product above). It is difficult to imagine what this braiding could mean. It is better to imagine braid and allow the strands to fuse and split (annihilation and pair creation vertices).

2. This braiding gives rise in the planar projection representation of braids to a generalization of non-planar Feynman diagrams. Non-planar diagrams are the basic unsolved problem of twistor approach and have prevented its development to a full theory allowing to construct exact expressions for the full scattering amplitudes (I remember however that Nima Arkani-Hamed et al have conjectured that non-planar amplitudes could be constructed by some procedure: they notice the role of permutation group and talk also about braidings (describable using covering groups of permutation groups)). In TGD framework the non-planar Feynman diagrams correspond to non-trivial braids for which the projection of braid to plane has crossing lines, say  $a$  and  $b$ , and one must decide whether the line  $a$  goes over  $b$  or vice versa.
3. An interesting open question is whether one must sum over all braidings or whether one can choose only single braiding. Choice of single braiding might be possible and reflect the failure of string

determinism for Kähler action and it would be favored by TGD as almost topological quantum field theory (TQFT) vision in which Kähler action for preferred extremal is topological invariant.

#### 4.8 Scattering amplitudes as generalized braid invariants

The last big idea is the reduction of quantum TGD to generalized knot/braid theory (I have talked also about TGD as almost TQFT). The scattering amplitude can be identified as a generalized braid invariant and could be constructed by the generalization of the recursive procedure transforming in a step-by-step manner given braided tree diagram to a non-braided tree diagram: essentially what Alexander the Great did for Gordian knot but tying the pieces together after cutting. At each step one must express amplitude as superposition of amplitudes associated with the different outcomes of splitting followed by reconnection. This procedure transforms braided tree diagram to a non-braided tree diagrams and the outcome is the scattering amplitude!

### References

- [1] Goncharov A. Volumes of hyperbolic manifolds and mixed Tate motives. <http://arxiv.org/abs/alg-geom/9601021>, 1996.
- [2] Adamo T. Twistor actions for gauge theory and gravity. Available at: <http://arxiv.org/pdf/1308.2820.pdf>, 2013.
- [3] Pitkänen M. Basic Extremals of Kähler Action. In *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdclass/tgdclass.html#class](http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#class), 2006.
- [4] Pitkänen M. Does Riemann Zeta Code for Generic Coupling Constant Evolution? In *Towards M-Matrix*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdquantum/tgdquantum.html#fermizeta](http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#fermizeta), 2006.
- [5] Pitkänen M. TGD and Cosmology. In *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdclass/tgdclass.html#cosmo](http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#cosmo), 2006.
- [6] Pitkänen M. The classical part of the twistor story. In *Towards M-Matrix*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdquantum/tgdquantum.html#twistorstory](http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#twistorstory), 2006.
- [7] Pitkänen M. Motives and Infinite Primes. In *TGD as a Generalized Number Theory*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdnumber/tgdnumber.html#infmotives](http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#infmotives), 2011.
- [8] Pitkänen M. Unified Number Theoretical Vision. In *TGD as a Generalized Number Theory*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdnumber/tgdnumber.html#numbervision](http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#numbervision), 2014.
- [9] Pitkänen M. What are the counterparts of Einstein's equations in TGD? In *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: [http://tgdtheory.fi/public\\_html/tgdclass/tgdclass.html#EinsteinTGD](http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#EinsteinTGD), 2013.
- [10] Pitkänen M. Cold Fusion Again . Available at: [http://tgdtheory.fi/public\\_html/articles/cfagain.pdf](http://tgdtheory.fi/public_html/articles/cfagain.pdf), 2015.
- [11] Pitkänen M. Does Riemann Zeta Code for Generic Coupling Constant Evolution? . Available at: [http://tgdtheory.fi/public\\_html/articles/fermizeta.pdf](http://tgdtheory.fi/public_html/articles/fermizeta.pdf), 2015.
- [12] Pitkänen M. Why the non-trivial zeros of Riemann zeta should reside at critical line? . Available at: [http://tgdtheory.fi/public\\_html/articles/rhagain.pdf](http://tgdtheory.fi/public_html/articles/rhagain.pdf), 2015.

- [13] Pitkänen M. Is Non-Associative Physics and Language Possible Only in Many-Sheeted Space-time?  
Available at: [http://tgdtheory.fi/public\\_html/articles/braidparse.pdf](http://tgdtheory.fi/public_html/articles/braidparse.pdf), 2015.